



Mathematics 2

المرحلة الأولى / المحاضرة الثانية

قسم الأنظمة الطبية الذكية

م. د. ايناس رضا علي

Matrices and Determinants

A matrix is a rectangular array of elements (scalars) from a field. The order, or size, of a matrix is specified by the number of rows and the number of columns, i.e. A an “ m by n ” matrix has m rows and n columns, and the element in the i th row and j th column is often denoted by a_{ij} :

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

A vector is a matrix with a single row (or column) of n elements , i.e. the column vector is:-

$$A = \begin{bmatrix} a_1 \\ a_2 \\ . \\ . \\ a_n \end{bmatrix} \quad \text{and row vector is } A = [a_1 \ a_2 \ . \ . \ a_n]$$

The matrix is square if the number of rows and columns are equal (i.e. $m = n$) and the elements a_{ij} of a square matrix are called the main diagonal.

$$\begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{bmatrix}$$

The identity matrix: $I =$ $\begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{bmatrix}$ **is square matrix**

with one in each main diagonal position and zeros else.

The diagonal matrix $D = \begin{bmatrix} a_1 & 0 & \dots & 0 \\ 0 & a_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_n \end{bmatrix}$ has the elements

a_1, a_2, \dots, a_n in its main diagonal position and zeros in all other locations, some of the a_i may be zero but not all.

A $n \times n$ triangular matrix has the pattern:-

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ 0 & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix}$$

lower triangular matrix

or

$$\begin{bmatrix} a_{11} & 0 & \dots & 0 \\ a_{21} & a_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

upper triangular matrix

The $m \times n$ null matrix:- $\theta = \begin{bmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 \end{bmatrix}$ has zero in each of its positions.

Elementary operations with matrices and vectors

1. Equality:- Two $m \times n$ matrices and A and B are said to be equal if: $a_{ij} = b_{ij} \quad \forall$ pairs of i and j .

EX-1 – Find the values of x, y for the following matrix equation:

$$\begin{bmatrix} x - 2y & 0 \\ -2 & 6 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ -2 & x + y \end{bmatrix}$$

Sol. –

$$\begin{array}{l} x - 2y = 3 \\ x - 2y = 3 \quad \dots(1) \\ x + y = 6 \quad \dots(2) * 2 \end{array} \Rightarrow \frac{2x + 2y = 12}{3x = 15} \Rightarrow \boxed{x = 5}$$

substitution $x = 5$ in (2) $\Rightarrow 5 + y = 6 \Rightarrow \boxed{y = 1}$

2. Addition:- The sum of two matrices of like dimensions is the matrix of the sum of the corresponding elements. If:-

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1n} \\ b_{21} & b_{22} & \dots & b_{2n} \\ \dots & \dots & \dots & \dots \\ b_{m1} & b_{m2} & \dots & b_{mn} \end{bmatrix}$$

then

$$A \mp B = \begin{bmatrix} a_{11} \mp b_{11} & a_{12} \mp b_{12} & \dots & a_{1n} \mp b_{1n} \\ a_{21} \mp b_{21} & a_{22} \mp b_{22} & \dots & a_{2n} \mp b_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} \mp b_{m1} & a_{m2} \mp b_{m2} & \dots & a_{mn} \mp b_{mn} \end{bmatrix}$$

thus:

- 1) $A+B = B+A$
- 2) $A+(B+C) = (A+B)+C$
- 3) $A-(B-C) = A-B+C$

EX-2- Find $A+B$ and $A-B$ if:-

$$A = \begin{bmatrix} 2 & 1 & 3 \\ 1 & 0 & -2 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & -2 & 2 \\ 2 & 3 & -1 \end{bmatrix}$$

Sol.-

$$A + B = \begin{bmatrix} 2+1 & 1-2 & 3+2 \\ 1+2 & 0+3 & -2-1 \end{bmatrix} = \begin{bmatrix} 3 & -1 & 5 \\ 3 & 3 & -3 \end{bmatrix}$$

$$A - B = \begin{bmatrix} 2-1 & 1-(-2) & 3-2 \\ 1-2 & 0-(+3) & -2-(-1) \end{bmatrix} = \begin{bmatrix} 1 & 3 & 1 \\ -1 & -3 & -1 \end{bmatrix}$$

3. Multiplication by a scalar:- The matrix A is multiplied by the scalar C by multiplying each element of A by c :-

$$CA = \begin{bmatrix} ca_{11} & ca_{12} & \dots & ca_{1n} \\ ca_{21} & ca_{22} & \dots & ca_{2n} \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ ca_{m1} & ca_{m2} & \dots & ca_{mn} \end{bmatrix}$$

EX-3- Assume $A = \begin{bmatrix} 3 & 2 & 1 \\ 0 & 5 & -1 \end{bmatrix}$, find $3A$.

Sol.-

$$3A = \begin{bmatrix} 3*3 & 3*2 & 3*1 \\ 3*0 & 3*5 & 3*(-1) \end{bmatrix} = \begin{bmatrix} 9 & 6 & 3 \\ 0 & 15 & -3 \end{bmatrix}$$

4. Matrix multiplication:- For the matrix product AB to be defined it is necessary that the number of columns of A be equal to the number of rows of B . The dimensions of such matrices are said to be conformable. If A is of dimensions $m \times p$ and B is $p \times n$, then the ij th element of the product $C=AB$ is computed as:-

$$C_{ij} = \sum_{k=1}^p a_{ik} b_{kj}$$

This is the sum of the products of corresponding elements in the i th row of A and j th column of B . The dimensions of AB are of course $m \times n$.

EX-4- Assume $A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 6 & 5 & 4 \\ -1 & 1 & -1 \\ 0 & 2 & 0 \end{bmatrix}$ find AB .

Sol.-

$$\begin{aligned} AB &= \begin{bmatrix} 1*6+2(-1)+3*0 & 1*5+2*1+3*2 & 1*4+2(-1)+3*0 \\ -1*6+0(-1)+1*0 & -1*5+0*1+1*2 & -1*4+0(-1)+1*0 \end{bmatrix} \\ &= \begin{bmatrix} 4 & 13 & 2 \\ -6 & -3 & -4 \end{bmatrix} \end{aligned}$$

Properties of multiplication:-

- a) $A(B+C) = AB + AC$ *distributive law*
- b) $A(BC) = (AB)C$ *associative law*
- c) $AB \neq BA$ *commutative law does not hold*
- d) $AI = IA = A$

EX-5- Assume $A = \begin{bmatrix} 1 & 2 \\ 0 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix}$, verify that $AB \neq BA$.

Sol.-

$$AB = \begin{bmatrix} 1 & 2 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 7 & 1 \\ 6 & 3 \end{bmatrix} \quad \& \quad BA = \begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 2 & 7 \end{bmatrix}$$

Hence $AB \neq BA$

5. Transpose of matrix:- Let A is any $m \times n$ matrix the transpose of A is $n \times m$ matrix A' formed by interchanging the role of rows and columns.

$$A' = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}' = \begin{bmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & \dots & a_{m2} \\ \dots & \dots & \dots & \dots \\ a_{1n} & a_{2n} & \dots & a_{mn} \end{bmatrix}$$

If a matrix is square and equal to its transpose, it is said to be symmetric, then $a_{ij} = a_{ji}$ for all pairs of i and j .

Properties of transpose are:-

- a) $(A \mp B)' = A' \mp B'$
- b) $(AB)' = B'A'$
- c) $(A')' = A$

EX-6- Assume $A = \begin{bmatrix} 3 & 2 & 5 \\ 2 & -1 & 4 \\ 5 & 4 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 4 & -1 & 0 \\ 5 & 4 & 3 \\ 2 & 1 & -1 \end{bmatrix}$, show that:-

- 1) A is symmetric matrix
- 2) $(A + B)' = A' + B'$
- 3) $(AB)' = B'A'$

Sol.-

$$1) A' = \begin{bmatrix} 3 & 2 & 5 \\ 2 & -1 & 4 \\ 5 & 4 & 0 \end{bmatrix}' = \begin{bmatrix} 3 & 2 & 5 \\ 2 & -1 & 4 \\ 5 & 4 & 0 \end{bmatrix} = A \Rightarrow A \text{ is a symmetric matrix.}$$

$$2) L.H.S. = (A+B)' = \begin{bmatrix} 7 & 1 & 5 \\ 7 & 3 & 7 \\ 7 & 5 & -1 \end{bmatrix}' = \begin{bmatrix} 7 & 7 & 7 \\ 1 & 3 & 5 \\ 5 & 7 & -1 \end{bmatrix}$$

$$R.H.S. = A' + B' = \begin{bmatrix} 3 & 2 & 5 \\ 2 & -1 & 4 \\ 5 & 4 & 0 \end{bmatrix} + \begin{bmatrix} 4 & 5 & 2 \\ -1 & 4 & 1 \\ 0 & 3 & -1 \end{bmatrix} = \begin{bmatrix} 7 & 7 & 7 \\ 1 & 3 & 5 \\ 5 & 7 & -1 \end{bmatrix} = L.H.S.$$

$$\therefore (A+B)' = A' + B'$$

$$3) L.H.S. = (AB)' = \begin{bmatrix} 32 & 10 & 1 \\ 11 & -2 & -7 \\ 40 & 11 & 12 \end{bmatrix}' = \begin{bmatrix} 32 & 11 & 40 \\ 10 & -2 & 11 \\ 1 & -7 & 12 \end{bmatrix}$$

$$R.H.S. = B'A' = \begin{bmatrix} 4 & 5 & 2 \\ -1 & 4 & 1 \\ 0 & 3 & -1 \end{bmatrix} \begin{bmatrix} 3 & 2 & 5 \\ 2 & -1 & 4 \\ 5 & 4 & 0 \end{bmatrix} = \begin{bmatrix} 32 & 11 & 40 \\ 10 & -2 & 11 \\ 1 & -7 & 12 \end{bmatrix} = L.H.S.$$

$$\therefore (AB)' = B'A'$$

6. Vector inner product:- The inner product of two vectors with the same number of elements is defined to be the sum of the products of the corresponding elements:-

$$A'B = [a_1 \quad a_2 \quad \dots \quad a_n] \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} = \sum_{i=1}^n a_i b_i$$

Since the inner product is a scalar, hence $A'B = B'A$. Moreover, the inner product of two vectors may be taken the following term:-

$$A B' = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} \begin{bmatrix} b_1 & b_2 & \cdots & b_n \end{bmatrix} = \begin{bmatrix} a_1 b_1 & a_1 b_2 & \cdots & a_1 b_n \\ a_2 b_1 & a_2 b_2 & \cdots & a_2 b_n \\ \cdots & \cdots & \cdots & \cdots \\ a_n b_1 & a_n b_2 & \cdots & a_n b_n \end{bmatrix}$$

Which is $n \times n$ matrix.

EX-7- Let $A = \begin{bmatrix} 5 \\ -2 \\ 1 \end{bmatrix}$ and $B = \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}$, find $A'B$ and AB'

Sol.-

$$A'B = [5 \quad -2 \quad 1] \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix} = 5 * 2 + (-2) * (-1) + 1 * 3 = 15$$

$$AB' = \begin{bmatrix} 5 \\ -2 \\ 1 \end{bmatrix} [2 \quad -1 \quad 3] = \begin{bmatrix} 10 & -5 & 15 \\ -4 & 2 & -6 \\ 2 & -1 & 3 \end{bmatrix}$$

Determinants

The minor of the element a_{ij} in a matrix A is the determinant of the matrix that remains when the row and column containing a_{ij} are deleted. For example, let:-

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \text{ then the minor of } a_{21} \text{ is } \begin{bmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{bmatrix}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix} \text{ then the minor of } a_{34} \text{ is } \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{41} & a_{42} & a_{43} \end{bmatrix}$$

and so on.

The cofactor of a_{ij} is the determinant A_{ij} that is $(-1)^{i+j}$ times the minor of a_{ij} . Thus:-

$$\text{for matrix } (3 \times 3) \Rightarrow A_{23} = (-1)^{2+3} \begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix} = - \begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix}$$

$$\text{for matrix } (4 \times 4) \Rightarrow A_{31} = (-1)^{3+1} \begin{vmatrix} a_{11} & a_{13} & a_{14} \\ a_{22} & a_{23} & a_{24} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} = \begin{vmatrix} a_{11} & a_{13} & a_{14} \\ a_{22} & a_{23} & a_{24} \\ a_{42} & a_{43} & a_{44} \end{vmatrix}$$

With each square matrix A we associate a number $\det A$ or $|A|$ or $|a_{ij}|$ called the determinant of A , calculated from the entries of A in the following way:-

$$\text{for } n=1, A = [a] \Rightarrow |A| = a$$

$$\text{for } n=2, A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \Rightarrow |A| = a_{11}a_{22} - a_{12}a_{21}$$

$$\text{for } n=3, A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \Rightarrow |A| = \begin{array}{r} a_{11} \quad a_{12} \quad a_{13} \\ a_{21} \quad a_{22} \quad a_{23} \\ a_{31} \quad a_{32} \quad a_{33} \end{array} \begin{array}{c} a_{11} \quad a_{12} \\ a_{21} \quad a_{22} \\ a_{31} \quad a_{32} \end{array} \begin{array}{c} a_{11} \quad a_{12} \\ a_{21} \quad a_{22} \\ a_{31} \quad a_{32} \end{array}$$

+ + +

$$|A| = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - (a_{13}a_{22}a_{31} + a_{11}a_{23}a_{32} + a_{12}a_{21}a_{33})$$

The determinant of a square matrix can be calculated from the cofactors of any row or any column.

EX-8- Find the determinant of the matrix:- $A = \begin{bmatrix} 2 & 1 & 3 \\ 3 & -1 & -2 \\ 2 & 3 & 1 \end{bmatrix}$

Sol.-

Ist method

$$A = \begin{bmatrix} 2 & 1 & 3 \\ 3 & -1 & -2 \\ 2 & 3 & 1 \end{bmatrix} \begin{array}{c} a_{11} \quad a_{12} \quad a_{13} \\ a_{21} \quad a_{22} \quad a_{23} \\ a_{31} \quad a_{32} \quad a_{33} \end{array} \begin{array}{c} a_{11} \quad a_{12} \\ a_{21} \quad a_{22} \\ a_{31} \quad a_{32} \end{array} \begin{array}{c} a_{11} \quad a_{12} \\ a_{21} \quad a_{22} \\ a_{31} \quad a_{32} \end{array}$$

+ + +

$$= 2(-1) \cdot 1 + 1(-2) \cdot 2 + 3 \cdot 3 \cdot 3 - (3(-1) \cdot 2 + 2(-2) \cdot 3 + 1 \cdot 3 \cdot 1)$$

$$= 36$$

2nd method

If we were to expand the determinant by cofactors according to elements of its third column, say, we would get:-

$$\begin{aligned}
 A &= a_{13}A_{13} + a_{23}A_{23} + a_{33}A_{33} \\
 &= 3(-1)^{1+3} \begin{vmatrix} 3 & -1 \\ 2 & 3 \end{vmatrix} + (-2)(-1)^{2+3} \begin{vmatrix} 2 & 1 \\ 2 & 3 \end{vmatrix} + 1(-1)^{3+3} \begin{vmatrix} 2 & 1 \\ 3 & -1 \end{vmatrix} \\
 &= 3(9 - (-2)) + 2(6 - 2) + (-2 - 3) = 36
 \end{aligned}$$

Useful facts about determinants:-

F-1: If two rows of matrix are identical, the determinant is zero.

EX-9 Show that:- $\begin{vmatrix} 3 & -1 & 2 \\ 2 & -3 & 5 \\ 3 & -1 & 2 \end{vmatrix} = 0$

Sol.-

$$\begin{vmatrix} 3 & -1 & 2 \\ 2 & -3 & 5 \\ 3 & -1 & 2 \end{vmatrix} \begin{vmatrix} 3 & -1 \\ 2 & -3 \\ 3 & -1 \end{vmatrix} = -18 - 15 - 4 - (-18 - 15 - 4) = 0$$

F-2: Interchanging two rows of matrix changes the sign of its determinants.

EX-10 Show that:- $\begin{vmatrix} 2 & 1 & 5 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{vmatrix} = -\begin{vmatrix} -1 & 0 & 3 \\ 2 & 1 & 5 \\ 1 & -2 & 4 \end{vmatrix}$

Sol.-

$$L.H.S. = \begin{vmatrix} 2 & 1 & 5 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{vmatrix} \begin{vmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -2 \end{vmatrix} = 0 + 3 + 10 - (0 - 12 - 4) = 29$$

$$R.H.S. = \begin{vmatrix} -1 & 0 & 3 \\ 2 & 1 & 5 \\ 1 & -2 & 4 \end{vmatrix} \begin{vmatrix} -1 & 0 \\ 2 & 1 \\ 1 & -2 \end{vmatrix} = -(-4 + 0 - 12 - (3 + 10 + 0)) = 29 = L.H.S.$$

$$\therefore \begin{vmatrix} 2 & 1 & 5 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{vmatrix} = -\begin{vmatrix} -1 & 0 & 3 \\ 2 & 1 & 5 \\ 1 & -2 & 4 \end{vmatrix}$$

F-3: The determinant of the transpose of a matrix is equal to the original determinant.

$$\underline{\text{EX-11}} \text{ Show that: } \begin{vmatrix} 2 & 1 & 5 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{vmatrix} = \begin{vmatrix} 2 & -1 & 1 \\ 1 & 0 & -2 \\ 5 & 3 & 4 \end{vmatrix}$$

Sol.-

$$L.H.S. = 29 \quad \text{from ex-10}$$

$$R.H.S. = \begin{vmatrix} 2 & -1 & 1 \\ 1 & 0 & -2 \\ 5 & 3 & 4 \end{vmatrix} \begin{matrix} |2 & -1 \\ 1 & 0 \\ 5 & 3| \\ = 0 + 10 + 3 - (0 - 12 - 4) = 29 \end{matrix} = L.H.S.$$

$$\therefore \begin{vmatrix} 2 & 1 & 5 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{vmatrix} = \begin{vmatrix} 2 & -1 & 1 \\ 1 & 0 & -2 \\ 5 & 3 & 4 \end{vmatrix}$$

F-4: If each element of same row (or column) of a matrix is multiplied by a constant C , the determinant is multiplied by C .

$$\underline{\text{EX-12}} \text{ Show that: } \begin{vmatrix} 6 & 3 & 15 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{vmatrix} = 3 \begin{vmatrix} 2 & 1 & 5 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{vmatrix}$$

Sol.-

$$L.H.S. = \begin{vmatrix} 6 & 3 & 15 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{vmatrix} \begin{matrix} |6 & 3 \\ -1 & 0 \\ 1 & -2| \\ = 0 + 9 + 30 - (0 - 36 - 12) = 87 \end{matrix}$$

$$R.H.S. = 3 * 29 = 87 = L.H.S.$$

$$\therefore \begin{vmatrix} 6 & 3 & 15 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{vmatrix} = 3 \begin{vmatrix} 2 & 1 & 5 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{vmatrix}$$

F-5: If all elements of a matrix above the main diagonal (or all below it) are zero, the determinant of the matrix is the product of the elements on the main diagonal.

$$\text{EX-13 Find: } \begin{vmatrix} 5 & 0 & 0 \\ 2 & 3 & 0 \\ 1 & -1 & 4 \end{vmatrix}$$

Sol.-

$$\begin{array}{|ccc|cc} 5 & 0 & 0 & 5 & 0 \\ 2 & 3 & 0 & 2 & 3 \\ 1 & -1 & 4 & 1 & -1 \end{array} = 60 + 0 + 0 - (0 - 0 - 0) = 60$$

$$\text{Or directly } 5 * 3 * 4 = 60$$

F-6: If each element of a row of a matrix is multiplied by a constant C and the results added to a different row, the determinant is not changed.

$$\text{EX-14 Show that } |A|=|B| \text{ if } A=\begin{bmatrix} 2 & 1 & 5 \\ -1 & 0 & 3 \\ 1 & -2 & 4 \end{bmatrix} \text{ and } B \text{ is the matrix}$$

resultant from multiplying row (1) by 2 and adding to row (3).

$$\text{i.e. } B=\begin{bmatrix} 2 & 1 & 5 \\ -1 & 0 & 3 \\ 5 & 0 & 14 \end{bmatrix}$$

Sol.-

$$|A|=29 \text{ from ex-10}$$

$$|B|=\begin{array}{|ccc|cc} 2 & 1 & 5 & 2 & 1 \\ -1 & 0 & 3 & -1 & 0 \\ 5 & 0 & 14 & 5 & 0 \end{array} = 0 + 15 + 0 - (0 - 0 - 14) = 29$$

$$\therefore |A|=|B|$$

$$\text{EX-15 Find } \begin{vmatrix} 1 & -2 & 3 & 1 \\ 2 & 1 & 0 & 2 \\ -1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \end{vmatrix}$$

Sol.-

$$\xrightarrow{-2R_1+R_2} \left| \begin{array}{cccc} 1 & -2 & 3 & 1 \\ 2 & 1 & 0 & 2 \\ -1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \end{array} \right| \quad \text{Expanding the determinant by using the first column.} \Rightarrow \left| \begin{array}{ccc|cc} 5 & 2 & 0 & 5 & 2 \\ 0 & 4 & -1 & 0 & 4 \\ 1 & -6 & 1 & 1 & -6 \end{array} \right| = 20 + 6 + 0 - (0 - 10 + 0) = 36$$

Linear Equations

There are many methods to solve a system of linear equations:

$$AX=B$$

I) **Row Reduction method** It is often possible to transform the linear equations step by step into an equivalent system of equations that is so simple it can be solved by inspection.

We start with $n \times (n+1)$ matrix $[A:B]$ whose first n columns are the columns of A and whose last column is B . We are going to transform this augmented matrix with a sequence of elementary row operations into $[I:S]$ where S is the solution of X .

$$2x + 3y - 4z = -3$$

EX-16 Solve the following linear equation: $x + 2y + 3z = 3$

$$3x - y - z = 6$$

Sol.

$$AX = B \quad \text{where} \quad A = \begin{bmatrix} 2 & 3 & -4 \\ 1 & 2 & 3 \\ 3 & -1 & -1 \end{bmatrix}, \quad X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, \quad B = \begin{bmatrix} -3 \\ 3 \\ 6 \end{bmatrix}$$

$$[A:B] = \left[\begin{array}{ccc|c} 2 & 3 & -4 & -3 \\ 1 & 2 & 3 & 3 \\ 3 & -1 & -1 & 6 \end{array} \right] \xrightarrow[-2R_2+R_1]{-3R_2+R_3} \left[\begin{array}{ccc|c} 0 & -1 & -10 & -9 \\ 1 & 2 & 3 & 3 \\ 0 & -7 & -10 & -3 \end{array} \right]$$

$$\xrightarrow[R_1 \text{ and } R_2]{\text{inter change}} \left[\begin{array}{ccc|c} 1 & 2 & 3 & 3 \\ 0 & -1 & -10 & -9 \\ 0 & -7 & -10 & -3 \end{array} \right] \xrightarrow[7R_2+R_3]{2R_2+R_1} \left[\begin{array}{ccc|c} 1 & 0 & -17 & -15 \\ 0 & -1 & -10 & -9 \\ 0 & 0 & 60 & 60 \end{array} \right]$$

$$\xrightarrow[\frac{1}{6}R_3+R_2]{\frac{17}{60}R_3+R_1} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & 60 & 60 \end{array} \right] \xrightarrow[R_2*(-1)]{R_3*\frac{1}{60}} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \end{array} \right] = [I:S]$$

$$\text{Hence } \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \boxed{x=2, y=-1, z=1}$$

II) Cramer's Rule When the determinant of the coefficient matrix A of the system $AX=B$ is not zero (i.e. $|A| \neq 0$) the system has a unique solution that it may be found from the formulas:

$X_i = \frac{|A_i|}{|A|}$ Where $|A_i|$ is the determinant of the matrix, comes from replacing the i th column in A by the column of constant B .

EX-17 Resolve example 16 using Cramer's rule:

Sol.

$$|A| = \begin{vmatrix} 2 & 3 & -4 \\ 1 & 2 & 3 \\ 3 & -1 & -1 \end{vmatrix} \begin{matrix} 2 & 3 \\ 1 & 2 \\ 3 & -1 \end{matrix} = -4 + 27 + 4 - (-24 - 6 - 3) = 60$$

$$|A_1| = \begin{vmatrix} -3 & 3 & -4 \\ 3 & 2 & 3 \\ 6 & -1 & -1 \end{vmatrix} \begin{matrix} -3 & 3 \\ 3 & 2 \\ 6 & -1 \end{matrix} = 6 + 54 + 12 - (-48 + 9 - 9) = 120$$

$$|A_2| = \begin{vmatrix} 2 & -3 & -4 \\ 1 & 3 & 3 \\ 3 & 6 & -1 \end{vmatrix} \begin{matrix} 2 & -3 \\ 1 & 3 \\ 3 & 6 \end{matrix} = -6 - 27 - 24 - (-36 + 36 + 3) = -60$$

$$|A_3| = \begin{vmatrix} 2 & 3 & -3 \\ 1 & 2 & 3 \\ 3 & -1 & 6 \end{vmatrix} \begin{matrix} 2 & 3 \\ 1 & 2 \\ 3 & -1 \end{matrix} = 24 + 27 + 3 - (-18 - 6 + 18) = 60$$

$$\therefore x = \frac{|A_1|}{|A|} = \frac{120}{60} \Rightarrow \boxed{x=2}$$

$$y = \frac{|A_2|}{|A|} = \frac{-60}{60} \Rightarrow \boxed{y=-1}$$

$$z = \frac{|A_3|}{|A|} = \frac{60}{60} \Rightarrow \boxed{z=1}$$

The same result in ex - 16.

Problems – 8

1) Let $A = \begin{bmatrix} 1 & 2 \\ 0 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 4 & -2 \end{bmatrix}$, $C = \begin{bmatrix} 3 & 1 \\ 4 & -1 \\ 0 & 2 \end{bmatrix}$,

$$D = \begin{bmatrix} 1 & 0 & 4 \\ 0 & 1 & 2 \\ 0 & -1 & 1 \end{bmatrix}, E = \begin{bmatrix} 3 & -1 \\ 4 & 2 \end{bmatrix}. \text{ Find:-}$$

$$\left. \begin{array}{lllll} a) AB & b) DC & c) (D+I)C & d) DC+C & e) DCB \\ f) EI & g) 3A+E & h) -5E+A & i) E(2B) \\ ans.: a) \begin{bmatrix} -1 & 10 & -1 \\ -4 & 16 & -8 \end{bmatrix} b) \begin{bmatrix} 3 & 9 \\ 4 & 3 \\ -4 & 3 \end{bmatrix} c,d) \begin{bmatrix} 6 & 10 \\ 8 & 2 \\ -4 & 5 \end{bmatrix} e) \begin{bmatrix} -6 & 42 & -9 \\ 1 & 20 & 6 \\ -7 & 4 & -18 \end{bmatrix} \\ f) \begin{bmatrix} 3 & -1 \\ 4 & 2 \end{bmatrix} g) \begin{bmatrix} 6 & 5 \\ 4 & 14 \end{bmatrix} h) \begin{bmatrix} -14 & 7 \\ -20 & -6 \end{bmatrix} i) \begin{bmatrix} 8 & 4 & 22 \\ 4 & 32 & 16 \end{bmatrix} \end{array} \right\}$$

2) Find the value of x :-

$$\begin{bmatrix} x & 4 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ 1 & 0 & 2 \\ 0 & 2 & 4 \end{bmatrix} \begin{bmatrix} x \\ -7 \\ \cancel{5/4} \end{bmatrix} = 0$$

$$\left(ans.: x = \frac{1}{2} \text{ or } x = 1 \right)$$

3) Find v and w if: $[5 \ v] = v[-2 \ 1]$.

$$\left(ans.: w = v = -\frac{5}{2} \right)$$

4) Let $A = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 2 \\ -1 & 3 \\ 5 & -2 \end{bmatrix}$, Find:-

$$a) 2A + B' \quad b) B'A' - I$$

$$\left(ans.: a) \begin{bmatrix} 2 & -3 & 9 \\ 2 & 5 & 6 \end{bmatrix} b) \begin{bmatrix} 10 & 19 \\ -5 & -6 \end{bmatrix} \right)$$

5) Let $A = \begin{bmatrix} 3 & 0 & 1 \\ 0 & 1 & 2 \\ -1 & 1 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 1 & -2 & 0 \\ -1 & 3 & 2 \end{bmatrix}$, Find:-

$(2A - I)B'$ and show that $(AB)' = B'A'$

$$\left(\text{ans.} : \begin{bmatrix} 5 & -1 \\ -2 & 11 \\ -6 & 26 \end{bmatrix} \right)$$

6) For what value of x will: $\begin{vmatrix} x & x & 1 \\ 2 & 0 & 5 \\ 6 & 7 & 1 \end{vmatrix} = 0$?
 $(\text{ans.}: x = 2)$

7) Let A be an arbitrary 3 by 3 matrix and let R_{12} be the matrix obtained from the 3 by 3 identity matrix by

interchanging row 1 and 2 : $R_{12} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. a) Compute $R_{12}A$

and show that you would get the same result by interchanging rows 1 and 2 of A . b) Compute AR_{12} and show that the result is that you would get by interchanging column 1 and 2 of A .

$$\left(\text{ans.}: a) \begin{bmatrix} a_{21} & a_{22} & a_{23} \\ a_{11} & a_{12} & a_{13} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} b) \begin{bmatrix} a_{12} & a_{11} & a_{13} \\ a_{22} & a_{21} & a_{23} \\ a_{32} & a_{31} & a_{33} \end{bmatrix} \right)$$

8) Solve the following determinants:-

a) $\begin{vmatrix} 2 & 3 & 1 \\ 4 & 5 & 2 \\ 1 & 2 & 3 \end{vmatrix}$ b) $\begin{vmatrix} 2 & -1 & -2 \\ -1 & 2 & 1 \\ 3 & 0 & -3 \end{vmatrix}$ c) $\begin{vmatrix} 2 & -1 & 3 \\ 1 & 0 & 2 \\ 0 & 2 & 1 \end{vmatrix}$

d) $\begin{vmatrix} 1 & 0 & -1 \\ 0 & 2 & -2 \\ 2 & 0 & 1 \end{vmatrix}$ e) $\begin{vmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & -2 & 1 \\ 0 & -1 & 0 & 7 \\ 3 & 0 & 2 & 1 \end{vmatrix}$ f) $\begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 2 \end{vmatrix}$

g) $\begin{vmatrix} 1 & -1 & 2 & 3 \\ 2 & -2 & 6 \\ 1 & 0 & 2 & 3 \\ -2 & 2 & 0 & -5 \end{vmatrix}$ h) $\begin{vmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{vmatrix}$

$$\left(\text{ans.}: a) -5 \quad b) 0 \quad c) -7 \quad d) 6 \\ e) 38 \quad f) 1 \quad g) 2 \quad h) -1 \right)$$

9) Solve the following system of equations:-

$$a) \ x + 8y = 4$$

$$3x - y = -13$$

$$b) \ 2x + 3y = 5$$

$$3x - y = 2$$

$$c) \ x + y + z = 2$$

$$2x - y + z = 0$$

$$x + 2y - z = 4$$

$$d) \ 2x + y - z = 2$$

$$x - y + z = 7$$

$$e) \ 2x - 4y = 6$$

$$x + y + z = 1$$

$$f) \ x - z = 3$$

$$2y - 2z = 2$$

$$2x + 2y + z = 4$$

$$5y + 7z = 10$$

$$2x + z = 3$$

$$g) \ x_1 + x_2 - x_3 + x_4 = 2$$

$$x_1 - x_2 + x_3 + x_4 = -1$$

$$x_1 + x_2 + x_3 - x_4 = 2$$

$$x_1 + x_3 + x_4 = -1$$

$$h) \ 2x - 3y + 4z = -19$$

$$6x + 4y - 2z = 8$$

$$x + 5y + 4z = 23$$

$$\left\{ \begin{array}{ll} \text{ans. : } a) x = -4, y = 1 & b) x = y = 1 \\ c) x = \frac{6}{7}, y = \frac{10}{7}, z = -\frac{2}{7} & d) x = 3, y = -2, z = 2 \\ e) x = 0, y = -\frac{3}{2}, z = \frac{5}{2} & f) x = 2, y = 0, z = -1 \\ g) x_1 = 2, x_2 = 0, x_3 = x_4 = -\frac{3}{2} & h) x = -2, y = 5, z = 0 \end{array} \right.$$