




DNA Replication

- Before each cell division, new copies must be made of each of the many molecules that form the cell, including the duplication of all DNA molecules. DNA replication is the name given to this duplication process, which enables an organism's genetic information its genes — to be passed to the two daughter cells created when a cell divides.
- DNA replication is the process of producing two identical copies from one original DNA molecule. This biological process occurs in all living organisms. It is the basis for biological inheritance. DNA is composed of two strands and each strand of the original DNA molecule serves as template for the production of the complementary strand, a process referred to as semiconservative replication.

After Watson and Crick proposed the double helix model of DNA, three models for DNA replication were proposed: conservative, semiconservative and dispersive.





DNA replication process

DNA replications process the process of DNA replication followed by proofreading or error-checking mechanisms to ensure correct reading of the genetic code, like all biological polymerization processes (Transcription and Translation, will be discussed later), the process involves 3 stages:

1- Initiation

2-Elongation

3- Termination


1- Initiation

The replication of both prokaryotic and eukaryotic DNAs starts at a unique sequence called the **origin of replication**, which serves as a specific binding site for proteins that initiate the replication process.


In E. coli, which has a single origin of replication on its one chromosome (as do most prokaryotes), it is approximately 245 base pairs long and is **rich in AT sequence** (rich in adenine and thymine bases), because A-T base pairs have two hydrogen bonds (rather than the three bonds in a C-G pair) and easy to break in this site.

The origin of replication (oriC) is recognized by certain proteins that bind to this site called Initiator proteins. These proteins (DnaA in prokaryotes, origin recognition complex in yeast) bind specifically to the AT-rich replicator sequence oriC to form a specific DnaA-oriC complex. An enzyme called helicase unwinds the DNA by breaking the hydrogen bonds between the nitrogenous base pairs. ATP hydrolysis is required for this process.

As the DNA opens up, Y-shaped structures called replication forks are formed. Two replication forks are formed at the origin of replication, and these get extended bi-directionally as replication proceeds. (two replication forks begin at a single replication origin in bacteria and proceed in opposite directions around the chromosome forming θ theta shape, which look like a bubble, moving away from the origin till reaching the opposite direction in one point called Ter Terminus (Teri)).



The mechanism of eukaryotic DNA replication is similar to that of prokaryotic DNA replication, but it is more complex. There are multiple origins of replication on the eukaryotic chromosome so multiple replication bubbles will form.



In yeast, which is a eukaryote, special sequences known as Autonomously Replicating Sequences (ARS) are found on the chromosomes. These are equivalent to the origin of replication in *E. coli*.




Single-strand binding proteins (SSBPs) bind to the single strands of DNA near the replication fork to prevent the ssDNA strands from winding back into a double helix, thus maintaining the strand separation.

One of the key players in DNA replication is the enzyme DNA polymerase, also known as DNA pol (there are many types of DNA polymerases in prokaryotes and eukaryotes will be discussed later).

DNA polymerase is able to add nucleotides only in the 5' to 3' direction (a new DNA strand can be only extended in this direction). It also requires a free 3'-OH group to which it can add nucleotides by forming a phosphodiester bond between the 3'-OH end and the 5' phosphate of the next nucleotide. This essentially means that it cannot add nucleotides if a free 3'-OH group is not available.

The problem is solved with the help of an RNA sequence that provides the free 3'-OH end. RNA primase synthesizes an RNA primer that is about five to ten nucleotides long and complementary to the DNA template.

Because this sequence primes the DNA synthesis, it is appropriately called the primer. DNA polymerase can now extend this RNA primer, adding nucleotides one by one that are complementary to the template strand. (example: A in the template strand is complement to T in new growing strand, and G in the template strand is complement to C in new growing strand) ... The primer is RNA rather than DNA because DNA polymerases cannot start chains de novo






2- Elongation step

DNA double helix is anti-parallel; that is, one strand is in the 5' to 3' direction and the other is oriented in the 3' to 5' direction. Both strands of parental DNA serve as templates for the synthesis of new DNA. A new DNA strand is always synthesized in a 5' to 3' direction. Thus, the replication of both the strands goes in two different ways.

One strand, which is complementary to the 3' to 5' parental DNA strand, is synthesized continuously in 5' to 3' direction towards the replication fork because the DNA polymerase III can add nucleotides in this direction. This continuously synthesized strand is known as the leading strand.

in prokaryote, DNA polymerase III begins the synthesis of the leading strand by using the RNA primer formed by primase (from 5' to 3' direction or the same direction as the replication fork movement) and add the nucleotides according to complement base-pairing to the template (A=T, T=A; G≡ C; C≡G)






However, all known DNA polymerases synthesize DNA in the 5'→3' direction but not in the 3' → 5' direction. How then does one of the daughter (lagging) DNA strands appear to grow in the 3'→5' direction?

The answer is

The other strand, complementary to the 5' to 3' parental DNA, is extended away from the replication fork discontinuously; in small fragments known as **Okazaki fragments**, each requiring a primer to start the synthesis (**this strand needs a new primer for each of the short Okazaki fragments**). Okazaki fragments are then synthesized via extension of these RNA primers by DNA polymerase. An important consequence of such RNA priming is that newly synthesized Okazaki fragments contain an RNA-DNA joint, the discovery of which provided critical evidence for the role of RNA primers in DNA replication. Okazaki fragments are named after the Japanese scientist Reiji Okazaki (1968) who first discovered them. The strand with the Okazaki fragments is known as the **lagging strand**.



Most current evidence indicates that DNA polymerases epsilon ϵ and Delta δ , respectively, perform the bulk of leading and lagging strand replication of the eukaryotic nuclear genome and Pol γ in the mitochondria).



As synthesis proceeds, the RNA primers are replaced by DNA. The primers are removed by the exonuclease activity of DNA polymerase I in prokaryotes, and the gaps are filled in by deoxyribonucleotides. The nicks that remain between the newly synthesized DNA (that replaced the RNA primer) and the previously synthesized DNA are sealed by the enzyme DNA ligase that catalyzes the formation of phosphodiester linkage between the 3'-OH end of one nucleotide and the 5' phosphate end of the other fragment; this is the reason why the synthesis of the lagging strand is more complicated than the leading strand.

A protein called the sliding clamp holds the DNA polymerase in place as it continues to add nucleotides. (sliding-clamp proteins and clamp-loading proteins) that load the polymerase onto the primer and maintain its stable association with the template

in eukaryote, the enzyme ribonuclease H (RNase H), instead of a DNA polymerase I, removes the RNA primer, which is then replaced with DNA nucleotides. The gaps that remain are sealed by DNA Ligase.

3-Termination Step:

1-Termination requires that the progress of the DNA replication fork must stop or be blocked. Termination at a specific locus, when it occurs, involves the interaction between two components: (1) a termination site sequence in the DNA, and (2) a protein which binds to this sequence to physically stop DNA replication. In various bacterial species, this is named the DNA replication terminus site-binding protein, or Ter protein.

2-Because bacteria have circular chromosomes, termination of replication occurs when the two replication forks meet each other on the opposite end of the parental chromosome. As a result, the replication forks are constrained to always meet within the termination region of the chromosome.

3-Removes the primer (RNA fragments), by 5'-3' exonuclease activity of polymerase I, and replaces the RNA nucleotides with DNA nucleotides. and fill the gaps.


4- When this is complete, a single nick on the leading strand and several nicks on the lagging strand can be found. Ligase works to fill these nicks in, thus completing the newly replicated DNA molecule.



6- Topoisomerase IV will: separate the two complete daughter chromosomes into two chromosomes. To form a continuous lagging strand of DNA, the RNA primers must eventually be removed from the Okazaki fragments and replaced with DNA.

Termination in Eukaryotic cell

- ❖ Eukaryote cells initiate DNA replication at multiple points in the chromosome, so replication forks meet and terminate at many points in the chromosome; these are not known to be regulated in any particular way. Because eukaryotes have linear chromosomes, DNA replication is unable to reach the very end of the chromosomes.
- ❖ Primer removal at the end of the chromosome leaves a gap that can't be filled in (there is no DNA polymerase coming along to fill in that piece. (Remember that DNA synthesis can ONLY occur 5'→3'). So, on every round of replication, a little piece is lost from the end of the chromosome.



Consequently, special mechanisms are required to replicate the terminal sequences of the linear chromosomes of eukaryotic cells. These terminal sequences (telomeres) consist of tandem repeats of simple-sequence DNA (short DNA sequences that are repeated over and over at the ends of the chromosomes). Short stretches are lost from telomeres at each round of replication. But that's alright because there is an enzyme called **TELOMERASE** that can refill the telomeres from an RNA template (which is



able to maintain telomeres by catalyzing their synthesis in the absence of a DNA template)

- ❖ Telomerase contains an RNA template to guide synthesis of new telomeric repeats

Remember:

Enzymes involved in DNA replication

- ❖ Primase: in fact, is RNA polymerase thus the formed primer is RNA rather than DNA and it will remove latter by DNA polymerase I
- ❖ Topoisomerase I: will break the 3' 5' phosphodiester bond converting super coiled to relax form which opposite to ligase. Relaxes the DNA from its super-coiled nature
- ❖ DNA Helicase Also known as helix destabilizing enzyme causes formation of Replication Fork due to broken hydrogen bonds
- ❖ DNA Gyrase (and Topoisomerase IV) ; this is a specific type of topoisomerase II convert relaxed form to super coiled.
- ❖ DNA Ligase Re-anneals the semi-conservative strands and joins Okazaki Fragments of the lagging strand.
- ❖ Telomerase Lengthens telomeric DNA by adding repetitive nucleotide sequences to the ends of eukaryotic chromosomes.
- ❖ DNA Polymerase Builds a new duplex DNA strand by adding nucleotides in the 5' to 3' direction. performs proof-reading and error correction.
- ❖ DNA clamp: A protein (unit from polymerase which prevents DNA polymerase III from dissociating from the DNA parent strand.
- ❖ Single-Strand Binding (SSB) Proteins Bind to ssDNA and prevent the DNA double helix from re-annealing after DNA helicase unwinds it thus maintaining the strand separation.

Types of DNA polymerases in Prokaryotes

Named in order of discovery, DNA polymerase I, II, III DNA polymerase III is the main polymerase for DNA replication (Main enzyme that adds nucleotides in the 5'-3'). DNA polymerase II is involved in DNA repair.



DNA Polymerase I has :

- 1- Proofreading exonuclease in ($3' \rightarrow 5'$ direction)
- 2- Primer removal exonuclease in ($5' \rightarrow 3'$) direction
- 3-DNA synthesis ($5' \rightarrow 3'$) replaces primer with newly synthesized DNA

Types of enzyme	Initiation activity	Polymerization $5' \rightarrow 3'$	Exonuclease activity $3' \rightarrow 5'$	Exonuclease activity $5' \rightarrow 3'$
DNA polymerase I	–	+	+	+
DNA polymerase II	–	+	+	–
DNA polymerase III	–	+	+	–

Types of DNA polymerase in Eukaryotic cell

The DNA polymerases of eukaryotes are in general less understandable than the DNA polymerases of prokaryotes. Eukaryotic cells have FIVE polymerases: four major nuclear DNA polymerases: DNA polymerase alpha (Pol α), DNA polymerase delta (Pol δ) and DNA polymerase epsilon (Pol ϵ), DNA polymerase beta (poly β), and one found in mitochondria: DNA polymerase Gamma (Poly γ).

1-Polymerase alpha (Pol α): it is the only enzyme has primase activity beside DNA polymerase, Pol α initiates DNA synthesis on both the leading and lagging strands by synthesizing a RNA/DNA hybrid primer.

Pol α is a heterotetramer composed of two primase subunits and two polymerase subunits. The primase subunits initiate DNA replication by synthesizing short (7–12 ribonucleotides) RNA primers, which are then extended by polymerase α .

2-Pol β Beta polymerase: excision repair and it is not highly active and is not very processive.


3-Pol γ Gamma polymerase: polymerization the mitochondrial DNA beside repairing by its exonuclease activity $3' \rightarrow 5'$

4-Pol δ delta and 5- ϵ epsilon polymerase: polymerization lagging (δ) and leading (ϵ) strand respectively $5' \rightarrow 3'$. In eukaryotes.



DNA can be synthesized in vitro by technique known as Polymerase Chain Reaction (PCR) it will be discussed in practical part of Molecular Biology course.

- DNA replication, the basis for biological inheritance, is a fundamental process occurring in all living organisms to copy their DNA.
- In the process of "replication" each strand of the original double-stranded DNA molecule serves as template for the reproduction of the complementary strand.
- Two identical DNA molecules have been produced from a single double-stranded DNA molecule.



- In a cell, DNA replication begins at specific locations in the genome, called "origins".
- Unwinding of DNA at the origin, and synthesis of new strands, forms a replication fork.
- In addition to DNA polymerase, the enzyme that synthesizes the new DNA by adding nucleotides matched to the template strand, a number of other proteins are associated with the fork and assist in the initiation and continuation of DNA synthesis.
- Cellular proofreading that ensure near perfect fidelity for DNA replication.

Questions Sheet:

1. During replication, Okazaki fragments elongate

- (a) leading strand towards the replication fork
- (b) lagging strand towards the replication fork



- (c) leading strand away from the replication fork
 - (d) lagging strand away from the replication fork
- 2. Which of the following enzymes separates the two strands of DNA during replication?**
- (a) Gyrase
 - (b) Topoisomerase
 - (c) Helicase
 - (d) DNA polymerase
- 3. DNA replication is**
- (a) conservative
 - (b) conservative and discontinuous
 - (c) semi-conservative and discontinuous
 - (d) semi-conservative and semi-discontinuous
- 4. Which of the following is used in DNA replication studies?**
- (a) *Neurospora crassa*
 - (b) *Drosophila melanogaster*
 - (c) *Escherichia coli*
 - (d) *Pneumococcus*
- 5. Which of the following helps in opening of DNA double helix in front of replication fork?**
- (a) topoisomerase
 - (b) DNA polymerase-I
 - (c) DNA gyrase
 - (d) DNA ligase
- 6. Termination of replication is triggered by**
- (a) DNA polymerase
 - (b) Helicase
 - (c) SSB
 - (d) Tus protein
- 7. DNA polymerase synthesizes**
- (a) DNA in 5'-3' direction
 - (b) DNA in 3'-5' direction
 - (c) mRNA in 3'-5' direction
 - (d) mRNA in 5'-3' direction
- 8. Association of histones H1 with nucleosome shows**
- (a) the occurrence of transcription
 - (b) the occurrence of replication
 - (c) exposed DNA double helix
 - (d) the condensation of DNA into chromatin fibre
- 9. The 3' – 5' phosphodiester linkage joins**
- (a) two DNA strands



-
- (b) two nucleotides
 - (c) a nitrogenous base with pentose sugar
 - (d) two nucleosides
- 10. The fragments of DNA are joined together by which of the following enzymes?**
- (a) Endonuclease
 - (b) DNA polymerase
 - (c) Primase
 - (d) Ligase