



جامعة المستقبل
AL MUSTAQBAL UNIVERSITY

كلية العلوم
قسم الفيزياء الطبية

Lecture: (3)

Subject: Types of heat in Biothermal physics

First Stage

Lecturer: 3



What is Heat?

Heat is the form of energy that is transferred between two materials of different temperature

This transfer of energy occurs because of **differences in the average**

.translational kinetic energy per molecule in the two materials

Heat flows from the material with higher temperature to the material with

lower temperature

The **SI** unit of heat is the joule, where $1 \text{ joule} = 1 \text{ newton} \times \text{meter}$

What is Temperature?

Temperature is a measure of average translational kinetic energy per molecule in a substance.

Temperature scales are **Fahrenheit, Celsius and Kelvin**



In Fahrenheit scale, water freezes at 32° and boils at 212° .

On the Celsius scale, water freezes at 0° and boils at 100° .

The scientific standard, is the Kelvin scale. 0 Kelvin is equal to -273.15° Celsius.

Converting Between Celsius, Kelvin, and Fahrenheit Scales

| To Convert From... | Use This Equation |
|-----------------------|---|
| Celsius to Fahrenheit | $T_{\circ F} = 9/5 T_{\circ C} + 32$ |
| Fahrenheit to Celsius | $T_{\circ C} = 5/9 (T_{\circ F} - 32)$ |
| Celsius to Kelvin | $T_K = T_{\circ C} + 273.15$ |
| Kelvin to Celsius | $T_{\circ C} = T_K - 273.15$ |
| Fahrenheit to Kelvin | $T_K = 5/9 (T_{\circ F} - 32) + 273.15$ |
| Kelvin to Fahrenheit | $T_{\circ F} = 9/5 (T_K - 273.15) + 32$ |



Types of heat:-

There are three types of heat:

1. Perceptible heat is the heat that can cause a change and a difference in the temperature of a substance

2. Latent heat, a specific amount of energy is required to change the solid form of a particular substances into a liquid or the liquid into a gas. It is energy required for change of state

3. Specific heat is the quantity of heat required to raise the temperature of one gram of a substance by one Celsius degree. The units of specific heat are usually calories or joules per gram per Celsius degree. example, the specific heat of water is 1 calorie (or 4.186 joules) per gram per Celsius degree.



$$c = \frac{\Delta E}{m\Delta\theta}$$
$$\Delta E = mc\Delta\theta$$

m = mass (kg)

c = specific heat capacity (J/kg°C)

ΔE = change in thermal energy (J)

$\Delta\theta$ = change in temperature (°C)

Example

A 250g copper pipe is heated from 10°C to 31°C. What is the energy needed to heat the pipe? The specific heat capacity of copper is 390 J/kg⁻¹°C⁻¹.

$$\Delta E = mc\Delta\theta$$

$$m = 250\text{g} \times 10^{-3} = 0.25\text{ kg}, \quad \Delta\theta = 31 - 10 = 21^\circ\text{C}$$

$$\Delta E = 0.25\text{ kg} \times 390\text{ J/kg}^{-1}\text{°C}^{-1} \times 21^\circ\text{C} = 2048\text{J}$$

- Q:**
1. What is 12.0 °C in kelvins?
 2. What is 32.0 °C in degrees Fahrenheit?
 3. What is used to measure temperature?
 4. what the difference between heat and temperature?