



Republic of Iraq
Ministry of Higher Education & Scientific research
Al-Mustaqbal University
Science College
Biochemistry Department

Introduction in Chemistry

For

First Year Student

Lecture 4

By

Dr. Karrar M. Obaid

2024-2025

Quantum Number

An atomic orbital is specified by four quantum numbers. These determine the orbital's size, shape, and orientation in the space of orbital.

1- **The principal quantum number (n)** tells the average relative distance of an electron from the nucleus, and the energy of the electron in an atom. (It can have only positive integer $n = 1, 2, 3, 4, \dots$). The larger the value of n , the higher energy and the larger orbital, or electron shell.

2- **The angular momentum quantum number (ℓ)** or orbital quantum number describes the shape of the orbital, and the shape is limited by the principal quantum number n (the shape of the region of space occupied by the electron), ($\ell = 0, 1, 2, 3, \dots, n-1$).

- **Notice;** the value of ℓ defines the shape of the orbital, and the value of n defines the size.
- **Notice;** Orbitals that have the same value of n but different values of ℓ are called subshells. These subshells are given different letters to help chemists distinguish them from each other.
- For example $\ell = 0, 1, 2, 3$
- **Name = s, p, d, f**

3- **The magnetic quantum number (m_ℓ)** represents the orientation of the region in space occupied by an electron with respect to an applied magnetic field. (The value of m_ℓ depends on the value of ℓ , $m_\ell = +1, 0, -1$).

4- **Spin Quantum Number(m_s):** of the spin axis of an electron, either clockwise or counter-clockwise. Only two values are allowed for $m_s +1/2$ and $-1/2$.

- **Notice;** the maximum number of electrons may the atomic subshell contains calculated by the relation $[2 \times (2\ell + 1)]$.

Subshells

The number of values of the orbital angular number ℓ can also be used to identify the number of subshells in a principal electron shell:

- When $n = 1$, $\ell = 0$ (ℓ takes on one value and thus there can only be one subshell)
- When $n = 2$, $\ell = 0, 1$ (ℓ takes on two values and thus there are two possible subshells)
- When $n = 3$, $\ell = 0, 1, 2$ (ℓ takes on three values and thus there are three possible subshells)
- After looking at the examples above, we see that the value of n is equal to the number of subshells in a principal electronic shell:

1. Principal shell with $n = 1$ has one s subshell ($\ell = 0$)
2. Principal shell with $n = 2$ has one s subshell and one p subshell ($\ell = 0, 1$)
3. Principal shell with $n = 3$ has one s subshell, one p subshell, and one d subshell ($\ell = 0, 1, 2$)

We can designate a principal quantum number, n , and a certain subshell by combining the value of n and the name of the subshell (which can be found using ℓ). For example, 3p refers to the third principal quantum number ($n=3$) and the p subshell ($\ell=1$).

n	l	m_l	m_s	Number of orbitals	Orbital Name	Number of electrons	Total Electrons
1 (K shell)	0	0	$\frac{1}{2}$ $-\frac{1}{2}$	1	1s	2	2
2 (L Shell)	0	0	$\frac{1}{2}$ $-\frac{1}{2}$	1	2s	2	8
	1	-1, 0, +1	$\frac{1}{2}$ $-\frac{1}{2}$	3	2p	6	
3 (M-shell)	0	0	$\frac{1}{2}$ $-\frac{1}{2}$	1	3s	2	18
	1	-1, 0, +1	$\frac{1}{2}$ $-\frac{1}{2}$	3	3p	6	
	2	-2, -1, 0, +1, +2	$\frac{1}{2}$ $-\frac{1}{2}$	5	3d	10	
4 (L-shell)	0	0	$\frac{1}{2}$ $-\frac{1}{2}$	1	4s	2	32
	1	-1, 0, +1	$\frac{1}{2}$ $-\frac{1}{2}$	3	4p	6	
	2	-2, -1, 0, +1, +2	$\frac{1}{2}$ $-\frac{1}{2}$	5	4d	10	
	3	-3, -2, -1, 0, +1, +2, +3	$\frac{1}{2}$ $-\frac{1}{2}$	7	4f	14	

Quantum Number Chart

Example: Give the name, magnetic quantum numbers, and number of orbitals for each sublevel with the given n and l quantum numbers: (a) $n = 3$, $l = 2$ (b) $n = 2$, $l = 0$ (c) $n = 5$, $l = 1$ (d) $n = 4$, $l = 3$ Solution:

	n	l	Subshell name	Possible m_l	No. of orbital
a	3	2	3d	+2, +1, 0, -1, -2	5
b	2	0	2s	0	1

c	5	1	5p	+1,0,-1	3
d	4	3	4f	+3,+2,+1,0,-1,-2,-3	7

Problem: What is wrong with each of the following quantum number designations and/or subshell names?

	n	ℓ	m ℓ	Name
a	1	1	0	1p
b	4	3	+1	4d
c	3	1	-2	3p


Solution:

(a) A sublevel of $n = 1$ can have only $\ell = 0$, not $\ell = 1$. The only possible subshell is 1s.


(b) A subshell with $\ell = 3$ is an f subshell, not a d subshell. The subshell name should be 4f.

c) A subshell with $\ell = 1$ can have only m ℓ of -1, 0, +1, not -2.

Example: Write a set of quantum numbers for the last electrons in ground state and ions in a Ne atom.



$$n = 2, \ell = 1, m\ell = -1, m_s = -1/2$$





***no ions for noble gas**


Last electron

Example: Give the value for all four quantum number for the last electron in the ground state of Cu_{29} .

solution



$$n = 3, \ell = 2, m\ell = -2, m_s = -1/2$$




Last electron

Atomic Orbital Shapes

Size of orbitals; defined as the surface that contains 90% of the total electron probability.

S Orbital

S orbitals are spherical with the nucleus at the center. When n increases, s orbitals become larger and higher in energy because of their increased distance from the nucleus. Note: Each subshell of the H atom has orbitals with a characteristic shape. s orbitals ($l = 0$) are spherically symmetrical around the nucleus.




P Orbitals

There are three P orbitals in each P subshell, occur in levels $n=2$ and greater. All P orbitals have the same basic shape (two lobes arranged along a straight line with the nucleus between the lobes)but differ in their orientations in space). We denote these orbitals as $2P_x$, $2P_y$, and $2P_z$. A $2P_x$ orbital has its greatest electron

probability along the x-axis, a 2Py orbital along the y-axis, and a 2Pz orbital along the z-axis.

Notice;

as the value of l increases, the number of orbitals in a given subshell increases, and the shapes of the orbitals become more complex. Energies of p orbitals: $2p < 3p < 4p$.



d Orbitals

Occur in levels $n = 3$ and greater. Two fundamental shapes; a. Four orbitals with four lobes for each one, centered in the plane indicated in the orbital label dxz , dyz , dxy , and $dx^2 - y^2$ b. Fifth orbital is uniquely shaped - two lobes along the z-axis and a belt centered in the xy plane dz^2 .

