



جامعة المستقبل
AL MUSTAQBAL UNIVERSITY
كلية العلوم
قسم الفيزياء الطبية


Anatomy

The Digestive System

Dr. Abdulhusein Mizhir Almaamuri


Introduction:

- **Digestive tract:** also called **alimentary tract** : includes mouth, esophagus, stomach, small intestine and large intestine.
- **Accessory organs** include: the teeth, tongue, and glandular organs such as salivary glands, liver, gallbladder, and pancreas.



Terms of Oral Cavity

The mouth or oral cavity is where the process of digestion begins




- **A. Vestibule:**
- **B. Oral cavity proper:**
- **C. Frenulum:**
 - 1. labial
 - 2. lingual
- **D. Teeth**
 - 1. incisors
 - 2. canines
 - 3. premolars
 - 4. molars
- **E. Gingiva**
- **F. Hard palate**
- **G. Soft palate**
- **H. Uvula**

• Teeth

(mechanical breakdown)

- Incisors used for cutting
- Canines used for stabbing and holding
- Molars large surface area used for grinding
- Primary or deciduous teeth 20
- Secondary or permanent teeth 32

The permanent dentition consists of 32 teeth.
four incisors,
two canines,
four premolars (or bicuspids),
four molars
two wisdom teeth (third molars) in each jaw.




Muscles of Mastication


- 1. masseter
- 2. temporalis
- 3. medial and lateral pterygoids.

Nerve supply: Mandibular nerve (CN V3)

The muscles of mastication are muscles that attach to the mandible and thereby produce movements of the lower jaw(temporomandibular joint).



Salivary Glands-three pairs



- **1. Parotid:** largest.
- **2. Sublingual:** smallest.
- **3. Submandibular:**
- Posterior half of inferior border of mandible.

Functions of Saliva:

1. Prevents bacterial infection
2. Lubrication
3. Contains salivary amylase that breaks down starch into disaccharides.
4. Helps to form **bolus** for **deglutition**

Histology of the Digestive Tract

1. Mucosa

2. Submucosa

3. Muscularis:

1. circular and longitudinal
2. smooth except for upper esophagus

4. Serosa or adventitia:


1. visceral peritoneum in abdominal cavity
2. tunica adventitia outside of the abdominal cavity

Nervous regulation of the Digestive System

– 1. Local: enteric nervous system

- Types of neurons: sensory, motor, interneurons
- Coordinates peristalsis and regulates local reflexes
- As stomach empties into small intestine, local reflex regulates rate of emptying

– 2. General: coordination with the CNS. May initiate reflexes because of sight, smell, or taste of food. Parasympathetic primarily. Sympathetic input inhibits muscle contraction, secretion, and decrease of blood flow to the digestive tract.



Pharynx

A. Common chamber of respiratory and digestive system


B. Structures ensuring that food gets to the right place

1. hard palate

2. soft palate and uvula

3. epiglottis: The function of the epiglottis is to close the laryngeal inlet during swallowing and so to prevent the passage of food and liquid into the lungs (aspiration). This is why we can't (and shouldn't try to) talk and breathe while swallowing


- Nasopharynx – not part of the digestive system
- Oropharynx – posterior to oral cavity
- Laryngopharynx – below the oropharynx and connected to the esophagus




(a)

Esophagus

- The **esophagus** is posterior to the larynx and trachea in the neck region and upper thorax. It passes through the diaphragm, and connects with the stomach. It's about 25 cm in length.
- There are inner circular and outer longitudinal muscle layers.
- The upper third is skeletal muscle (voluntary), middle third is mixed, and lower third is smooth muscle (involuntary).
- Conducts food by peristalsis (slow rhythmic squeezing):
 - contraction of circular layer above the food and contraction of longitudinal below the food
- **Esophagogastric junction** is located approximately at the level of the diaphragm. Contractions of the diaphragm create sphincter-like effects, preventing reflux of stomach acids and content. The esophagogastric junction is a functional, not anatomical, sphincter.



Peristalsis in Esophagus



Stomach

The stomach is a muscular organ located on the left side of the upper abdomen. The stomach receives food from the oesophagus. As food reaches the end of the oesophagus, it enters the stomach through a muscular valve called the lower oesophageal sphincter.

The stomach is supplied with arterial blood from a branch of the celiac artery and venous blood leaves the stomach via the hepatic vein. The vagus nerve innervates the stomach with parasympathetic fibres that stimulate gastric motility and the secretion of gastric juice. Sympathetic fibres from the celiac plexus reduce gastric activity.

The stomach has the same four layers of tissue as the digestive tract but with some differences. The muscularis contains three layers of smooth muscle instead of two. It has longitudinal, circular and oblique muscle fibres. The extra muscle layer facilitates the churning, mixing and mechanical breakdown of food. The stomach secretes acid and enzymes that digest food. Ridges of muscle tissue called rugae line the stomach.

The pyloric sphincter is a muscular valve that opens to allow food to pass from the stomach to the duodenum.

Stomach

1. Openings


- **Gastroesophageal (cardiac)**: to esophagus
- **Pyloric**: to duodenum

2. Parts

- **Cardiac**
- **Fundus**
- **Body**
- **Pyloric: antrum and canal**
- **Greater and lesser curvatures**: attachment sites for omenta

1. Layers (from outside)

- a. Serosa or visceral peritoneum
- b. Muscularis: three layers
 - Outer longitudinal
 - Middle circular
 - Inner oblique
- c. Submucosa
- d. Mucosa
- **e. Rugae**: folds in mucosa



Peritoneum:



is the largest serous membrane of the body. Divide into:

1. **Parietal peritoneum:** lines the wall of abdominopelvic cavity internally
2. **Visceral peritoneum:** cover some of the organs in the cavity


The space between them contains fluid and is called **peritoneal cavity**. This cavity may be accumulated by several liters of fluid in a state called **ascites**.

Retroperitoneal organs: e.g., kidneys, pancreas, duodenum

Mesenteries: double sheets of peritoneum, surrounding and suspending portions of the digestive organs



Small Intestine




Anterior view

- Site of greatest amount of digestion and absorption of nutrients and water
- Divisions
 - **Duodenum**- first 25 cm beyond the pyloric sphincter.
 - **Jejunum**- 2.5 m
 - **Ileum**- 3.5 m. Peyer's patches or lymph nodules

Modifications to Increase Surface Area


- **1. Plicae circulares** (circular folds)
- **2. Villi** that contain capillaries and lacteals. Folds of the mucosa
- **3. Microvilli**: folds of cell membranes of absorptive cells
- 4. total surface area about that of a tennis court.

Ileocecal junction: where ileum meets large intestine. **Ileocecal sphincter** and **ileocecal valve**



Large Intestine

- A. Extends from ileocecal junction to **anus**
- B. Consists of **cecum**, **colon**, **rectum**, **anal canal**
- C. Movements sluggish (18-24 hours); chyme converted to feces.
- D. Absorption of water and salts, secretion of mucus, extensive action of microorganisms.
- E. 1500 mL chyme enter the cecum, 90% of volume reabsorbed yielding 80-150 mL of feces




Liver

The liver is the largest solid organ in the body. In adults, the liver can weigh up to 1.5 kilograms (kg). It is in the upper-right abdomen, just under the rib cage and below the diaphragm.

Hepatic Portal Circulation

The **hepatic artery** comes off the celiac trunk which in turn comes from the aorta. The venous blood from the digestive tract is collected by the **portal vein**, which then supplies blood to liver.

The **hepatic veins** drain blood from liver into the **inferior vena cava**. Branches of the hepatic **artery** and **vein** and the **bile duct** flow into the liver. Collectively, these three vessels are termed the **portal triad**.



Pancreas:

Retroperitoneal :compose of head, body and tail

Endocrine and exocrine gland

Endocrine: pancreatic islets or Islets of Langerhans.


Gall Bladder :

Sac where bile arrives constantly from liver, stored and concentrated.

Bile exits through cystic duct then into common bile duct

Gallstones:

Can block cystic duct





THANK YOU!

