

PRESCRIPTION WRITING

part 2

PRESCRIPTION

NAME: _____

ADDRESS: _____

Rx

SIGNATURE: X _____



UTI

Rx

**1-Trimethoprim-Sulfamethoxazole
400/80 mg tab**

**2 tablets every 12 hours
For 3 days**

**2-Ibuprofen 200 mg tab
1 tablet three times daily after meal**



Rx

**1-Cefixime 400 mg cap
1 capsule daily
For 5 days**

**2-Mefenamic acid (ponstan) 500
mg tab
1 tablet on need after meal**

Rx

**1-Ciprofloxacin 500 mg tab
1 tablet every 12 hours
For 7 days**

**2-Mefenamic acid (ponstan) 500
mg tab
1 tablet on need after meal**

Tonsillitis

Rx

1-Azithromycin 500 mg tab
1 tablet daily (1 hour before or
2 hours after a meal)
For 3 days

2-Mefenamic acid (ponstan) 250
mg cap
1 capsule TID after meal



Rx

1-Augmentin 625 mg cap
1 capsule every 8 hours
For 5 days

2-Paracetamol 500 mg tablet
2 tablet TID after meal

Children (Ex, 6 years old)

Rx

1-Azithromycin 200 mg/5 ml
suspension
5 ml once daily
For 3 days

2-Paracetamol 250 mg/5ml susp
5 ml QID

Osteoarthritis

Rx

**Meloxicam 15 mg tab
1 tab daily after meal**



Rx

**Diclofenac sodium (voltaren) 50 mg tab
1 tab twice daily after meal**

GI infection (common food poisoning)

Rx

Metronidazole (Flagyl) 500 mg tab

1 tab every 8 hours

For 3 days



Rx

Trimethoprim-Sulfamethoxazole 400/80 mg tab

2 tablets every 12 hours

For 3 days

Children (Ex, 6 years old)

Rx

Metronidazole (Flagyl) 200mg/5ml susp

5 ml every 8 hours

For 3 days

Common antibiotic injections

❖ **Ceftriaxone** 250 mg ; 500 mg ; 1 g IM or IV

Given twice daily

Used for many infections like respiratory tract infections, urinary tract infections and skin infections.

❖ **Cefotaxime (Claforan)** 250 mg ; 500 mg ; 1 g IM or IV

Given three times daily

Used for many infections like GIT infections, urinary tract infections and skin infections.



Severe renal infection, fever, pain, N & V


Rx

**Ceftriaxone 1 g vial IM
1 vial twice daily given IM
For 5 days**

**Diclofenac Na (Voltaren) 75 mg ampoule IM
1 amp daily**



Oral Infections Treatment



As a dentist you should avoid developing oral infections

Oral infections have a few common causes:

- **An infection of the pulp (center) of a tooth**
- **An infection from a wisdom tooth**
- **An infection that occurs after a tooth is removed**
- **A gum infection**
- **An infection caused by an injury**
- **An infection caused by blocked flow of saliva**

❖ *Symptoms of an infection include:*

- ***Pain***
- ***Swelling***
- ***Difficulty opening the mouth***
- ***Redness***
- ***Fever***
- ***Problems swallowing or breathing (with severe infections).***



Rx

**1- Erythromycin 250 mg cap
1 cap every 6 hours one hour before meal
For 5 days**

**2- Metronidazole (Flagyl) 500 mg tab
1 tab every 8 hours
For 5 days**

**3- Mefenamic acid (ponstan) 250 mg cap
1 capsule TID after meal**



Rx

1) Rodogyl tab (750,000 IU spiramycin + Metronidazole 125 mg)

2 tablets every 8 hours

2) Ibuprofen 200 mg tab

1 tablet three times daily after meal

Child 6 years or if < 6 the dose according to weight

Rx


Co-amoxiclav 457mg/5ml syrup

5 ml twice daily

For 5 days

Ibuprofen 100mg/5ml syrup

5 ml twice daily on need



Rx

Erythromycin 125mg/5ml susp

5ml every 6 hours

For 5 days

Mefenamic acid 50 mg/5ml syrup

5 ml on twice daily



12 years old patient

Rx

**1) Rodogyl tab (750,000 IU spiramycin +
Metronidazole 125 mg)**

1 tab TID

2) Ibuprofen 100mg/5ml syrup

5 ml TID



***Rodogyl :Children from 6 -10 years 1 tab BID, 10-15 years 1
tab TID***



Tips for Teeth Pain Management

Here are a few tips to help minimize your discomfort and speed recovery:

- ***we can put ice packs on the face to reduce swelling.***
- ***Eat soft and cool foods for a few days.***
- ***Wash the mouth with warm salt water. Use one-half teaspoon of salt in a cup of water.***
- ***Reduce smoking.***



Indications to use Mouth washes

- ☐ **Cases of excessive formation of plaque.**
- ☐ **Prevention in the treatment of gingivitis.**
- ☐ **People with difficulty in performing normal dental hygiene.**
- ☐ **Mouth wash before and after surgery.**
- ☐ **Prevention of dental caries.**

Ex. Chlorhexidine mouth wash.





Thank You