

General Physics

Lecture 6 / Theoretical



Standard Units of Measurement

First stage

Dr. Ahmed Najm Obaid

Assis. Lect. Amin Kadhum Awad

2024-2025

General Physics

Measurement

Physics is the study of interactions of matter and energy in all their diverse forms. Similar to all scientists, physicists strive for exactness or certainty in describing these interactions. They try to remove the uncertainties by eliminating subjective descriptions of events. Assuming that all measurements are correctly made, all observers who use the methods of physics will obtain exactly the same results. In addition to seeking certainty, physicists strive for simplicity; therefore, only three measurable quantities are considered basic. These base quantities are mass, length, and time, and they are the building blocks of all other quantities.

The secondary quantities are called derived quantities because they are derived from a combination of one or more of the three base quantities. For example, volume is length cubed (l^3), mass density is mass divided by volume (m/l^3), and velocity is length divided by time (l/t).

Units

Every measurement has two parts: a magnitude and a unit. For example, the SID is 100 cm. The magnitude, 100, is not meaningful unless a unit is also designated. Here, the unit of measurement is the centimeter.

SI Prefixes

Factor	Prefix	Symbol
1018	Exa	E
10 ¹⁵	Peta	Р
1012	Tera	T
10^9	Giga	G
10^{6}	Mega	M
10^{3}	Kilo	k
10^{2}	Hecto	h
10 ¹	Deca	da
10^{-1}	Deci	d
10^{-2}	Centi	С
10^{-3}	Milli	m
10^{-6}	Micro	μ
10^{-9}	Nano	n
10^{-12}	Pico	р
10^{-15}	Femto	р f
10^{-18}	Atto	a

SI Base Units

Name	Symbol
Meter	m
Kilogram	kg
Second	S
Ampere	Α
	Meter Kilogram Second

Special Quantities of Radiologic Science and Their Associated Special Units

	CUSTOMARY UNIT			SI UNIT	
Quantity	Name		Symbol	Name	Symbol
Exposure	roentgen		R	air kerma	Gy _a
Absorbed dose	rad		rad	gray	Gy_1
Effective dose	rem		rem	seivert	Sv
Radioactivity	curie		Ci	becquerel	Bq
Multiply	R	by	0.01	to obtain	Gya
Multiply	rad Gy	by	0.01	to obtain	Gy_t
Multiply	rem	by	0.01	to obtain	Sv
Multiply	Ci	by	3.73×10^{10}	to obtain	Bq
Multiply	R	by	2.583×10^{-4}	to obtain	C/kg

Question: The dimensions of a box are 30 cm \times 86 cm

 \times 4.2 m. Find the volume.

Answer: Formula for the volume of an object:

 $V = length \times width \times height or V = lwh$

Because the dimensions are given in different systems of units, however, we must choose only one system. Therefore,

$$V = (0.3 \text{ m}) (0.86 \text{ m}) (4.2 \text{ m})$$

= 1.1 m³

Question: Find the mass density of a solid box 10 cm

on each side with a mass of 0.4 kg.

Answer: D = mass/volume (change 10 cm to 0.1 m)

 $= 0.4 \text{ kg/}(0.1 \text{ m} \times 0.1 \text{ m} \times 0.1 \text{ m})$

 $= 0.4 \text{ kg}/0.001 \text{ m}^3$

 $= 400 \text{ kg/m}^3$

Mechanics

Mechanics is a segment of physics that deals with objects at rest (statics) and objects in motion (dynamics).

Velocity

The motion of an object can be described with the use of two terms: velocity and acceleration. Velocity, sometimes called speed, is a measure of how fast something is moving or, more precisely, the rate of change of its position with time. The velocity of a car is measured in kilometers per hour (miles per hour). Units of velocity in SI are meters per second (m/s). The equation for velocity (v) is as follows



Velocity

$$v = \frac{d}{t}$$

where *d* represents the distance traveled in time *t*.

The velocity of light is constant and is symbolized by c: $c = 3 \times 10^8$ m/s.

Question: What is the velocity of a ball that travels

Answer:
$$v = \frac{d}{t}$$

$$v = 60 \text{ m/4 s},$$


 $v = 15 \text{ m/s}$

Answer:
$$v = \frac{d}{d}$$

$$= \frac{6.69 \times 10^8 \ mi}{hr} \times \frac{1609 \ m/mi}{3600 \ s/hr}$$
$$= 2.99 \times 10^8 \ m/s$$


Acceleration

The rate of change of velocity with time is acceleration. It is how "quickly or slowly" the velocity is changing. Because acceleration is velocity divided by time, the unit is meters per second squared (m/s²). If velocity is constant, acceleration is zero. On the other hand, a constant acceleration of 2 m/s² means that the velocity of an object increased by 2 m/s each second. The defining equation for acceleration is given by the following:



Question: What is the acceleration of the dragster?

Answer: $a = \frac{80 \text{ m/s} - 0 \text{ m/s}}{10.2 \text{ s}}$ = 7.8 m/s²



 $v_o = 0 \text{ m/s} \quad a = 7.8 \text{ m/s}^2 \quad v_f = 80 \text{ m/s} \quad t = 10.2 \text{ s}$

Comprehensive Set of MCQs

1. What is the formula for calculating velocity?

- A) Distance × Time
- B) Distance / Time
- C) Time / Distance
- D) Distance + Time
- E) Distance Time

2. Which of the following is a base quantity?

- A) Force
- B) Energy
- C) Mass
- D) Velocity
- E) Acceleration

3. What is the SI unit of length?

- A) Kilogram
- B) Meter
- C) Liter
- D) Newton
- E) Joule

4. Which equation represents the relationship between mass, density, and volume?

- A) Density = Mass + Volume
- B) Density = Mass \times Volume
- C) Density = Mass / Volume
- D) Density = Volume / Mass
- E) Density = Mass Volume

5. What does the term 'acceleration' refer to?

- A) The speed of an object
- B) The rate of change of velocity
- C) The distance traveled
- D) The mass of an object
- E) The force applied

6. Which of the following describes the concept of velocity?

- A) Distance traveled
- B) Rate of change of position
- C) Total distance
- D) Average speed
- E) Instantaneous speed

7. What is the unit of measurement for time in the SI system?

- A) Second
- B) Minute
- C) Hour
- D) Day
- E) Year

8. Which of the following is NOT a derived quantity?

- A) Velocity
- B) Acceleration
- C) Mass
- D) Force
- E) Pressure

9. What is the formula for calculating force?

- A) Mass × Acceleration
- B) Mass / Acceleration
- C) Mass + Acceleration
- D) Mass Acceleration
- E) Acceleration / Mass

10. Which of the following describes mechanics?

- A) The study of light
- B) The study of sound
- C) The study of objects at rest and in motion
- D) The study of chemical reactions
- E) The study of electricity

11. What is the formula for calculating kinetic energy?

- A) $KE = 1/2 \text{ mv}^2$
- B) $KE = mv^2$
- C) KE = mgh
- D) KE = 1/2 mgh
- E) KE = mv

12. What is the unit of force in the SI system?

- A) Joule
- B) Newton
- C) Pascal
- D) Watt
- E) Coulomb

13. Which of the following is a vector quantity?

- A) Speed
- B) Distance
- C) Mass
- D) Displacement
- E) Temperature

14. Which of the following describes acceleration?

- A) Change in velocity over time
- B) Change in distance over time
- C) Change in mass over time
- D) Change in force over time
- E) Change in energy over time

15. Which of the following is a unit of energy?

- A) Joule
- B) Newton
- C) Pascal
- D) Meter
- E) Kilogram