


### 3. Vectors analysis/angle between two planes:-

*Angle between planes*

The angle between two intersecting planes is defined to be the acute angle between their normal vectors.

$$\theta = \cos^{-1} \left( \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1| |\mathbf{n}_2|} \right)$$



**Example:** Find the angle between the planes  $2x - 6y - z = 5$  and  $x + 2y - 2z = 12$

**Solution//**

$$\mathbf{n}_1 = 2\mathbf{i} - 6\mathbf{j} - \mathbf{k}$$

$$\mathbf{n}_2 = \mathbf{i} + 2\mathbf{j} - 2\mathbf{k}$$

$$|\mathbf{n}_1| = \sqrt{4 + 36 + 1} = \sqrt{41}$$

$$|n2| = \sqrt{1+4+4} = \sqrt{9}$$

$$\theta = \cos^{-1} \frac{\mathbf{n1} \cdot \mathbf{n2}}{|\mathbf{n1}| \cdot |\mathbf{n2}|} = \cos^{-1} \frac{-8}{\sqrt{41} \cdot \sqrt{9}} = 114.6^\circ$$

#### 4. Vectors analysis/intersection line & plane:-

**Example:** Find the vector parallel to the line of intersection of the planes  $3x-6y-2z=15$ ,  $x+2y-z=5$ .

**Solution/**

$$\mathbf{N1} = 3\mathbf{i} - 6\mathbf{j} - 2\mathbf{k}$$

$$\mathbf{N2} = \mathbf{i} + 2\mathbf{j} - \mathbf{k}$$

$$\mathbf{N} = \mathbf{N1} \times \mathbf{N2} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -6 & -2 \\ 1 & 2 & -1 \end{vmatrix}$$

$$= 10 \mathbf{i} + \mathbf{j} + 12\mathbf{k}$$

#### 5. Vector Functions:-

A vector –valued function of real variable can be written in component form as:

$$\mathbf{F}(t) = F_1(t)\mathbf{i} + F_2(t)\mathbf{j} + F_3(t)\mathbf{k}$$

#### 1. Limits

If  $\mathbf{L} = L_1\mathbf{i} + L_2\mathbf{j} + L_3\mathbf{k}$  is a vector in space

$\mathbf{F}(t)$  is a vector function

$$\mathbf{F}(t) = f(t)\mathbf{i} + g(t)\mathbf{j} + h(t)\mathbf{k}$$

$$\lim_{t \rightarrow a} f(t) = \lim_{t \rightarrow a} f_1(t) + \lim_{t \rightarrow a} f_2(t) + \lim_{t \rightarrow a} f_3(t)$$

**Example:** Find  $\lim_{t \rightarrow \pi} f(t)$  If  $f(t) = \cos t \mathbf{i} + 3 \sin t \mathbf{j} + t^3 \mathbf{k}$

**Solution//**

$$\begin{aligned}\lim_{t \rightarrow \pi} f(t) &= \lim_{t \rightarrow \pi} (\cos t \mathbf{i} + 3 \sin t \mathbf{j} + t^3 \mathbf{k}) \\ &= \lim_{t \rightarrow \pi} \cos t \mathbf{i} + \lim_{t \rightarrow \pi} 3 \sin t \mathbf{j} + \lim_{t \rightarrow \pi} t^3 \mathbf{k} = -\mathbf{i} + 0\mathbf{j} + \pi^3 \mathbf{k}\end{aligned}$$

## 2.Derivative.

$$\mathbf{r}(t) = \mathbf{f}(t)\mathbf{i} + \mathbf{g}(t)\mathbf{j} + \mathbf{h}(t)\mathbf{k}$$

$$\Delta \mathbf{r} = \mathbf{r}(t + \Delta t) - \mathbf{r}(t) \dots \dots \dots 1$$

Then if  $\mathbf{r}(t)$  sub in equation 1

$$\begin{aligned}\Delta \mathbf{r} &= \{\mathbf{f}(t + \Delta t) - \mathbf{f}(t)\}\mathbf{i} + \{\mathbf{g}(t + \Delta t) - \mathbf{g}(t)\}\mathbf{j} \\ &\quad + \{\mathbf{h}(t + \Delta t) - \mathbf{h}(t)\}\mathbf{k}\end{aligned}$$

As  $\Delta t = 0$

$$\begin{aligned}\frac{d\mathbf{r}}{dt} &= \lim_{\Delta t \rightarrow 0} \frac{\Delta \mathbf{r}}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} \frac{\{\mathbf{g}(t + \Delta t) - \mathbf{g}(t)\}\mathbf{j}}{\Delta t} + \lim_{\Delta t \rightarrow 0} \frac{\{\mathbf{h}(t + \Delta t) - \mathbf{h}(t)\}\mathbf{k}}{\Delta t} + \lim_{\Delta t \rightarrow 0} \frac{\{\mathbf{f}(t + \Delta t) - \mathbf{f}(t)\}\mathbf{i}}{\Delta t}\end{aligned}$$

$$\frac{d\mathbf{r}}{dt} = \frac{df}{dt} \mathbf{i} + \frac{dg}{dt} \mathbf{j} + \frac{dh}{dt} \mathbf{k}$$

**Notes//**

$$1. \text{ Velocity} = \frac{dr}{dt} = \bar{V}$$

$$2. \text{ Acceleration } a = \frac{d^2r}{dt^2} = \frac{dv}{dt}$$

$$3. \text{ Speed or magnitude of velocity} = |V|$$

Or velocity  $\bar{V} = \text{speed}|V| * \text{direction}$

**Example:** Find speed and direction of  $r(t)$  when  $t=2$     If  $r(t) = t^2\mathbf{i} + 2t^2\mathbf{j} + 5\mathbf{k}$

**Solution//**

$$\frac{dr}{dt} = \bar{V} = 3t^2\mathbf{i} + 4t\mathbf{j} + 0\mathbf{k}$$

$$\text{speed} = |V| = \sqrt{(3t^2)^2 + (4t)^2}$$

$$\text{At } t=2 \rightarrow |V| = 14.4$$

$$\text{Direction (at } t=2) = \frac{\bar{v}}{|V|}$$

$$= \frac{12\mathbf{i} + 8\mathbf{j} + 0\mathbf{k}}{14.4}$$

**Differential rules: -**

$$1. \frac{dc}{dt} = 0 \text{ if } c = \text{constant}$$

$$\text{Example : } c = 2\mathbf{i} + 4\mathbf{j} + 5\mathbf{k}, \frac{dc}{dt} = 0\mathbf{i} + 0\mathbf{j} + 0\mathbf{k} = \mathbf{0}$$

$$2. \text{ if } u(t) \text{ is a vector function, then } \frac{d cu}{dt} = c \cdot \frac{du}{dt}$$

**where c is constant a vector**

$$3. \frac{d(u \pm v)}{dt} = \frac{du}{dt} \pm \frac{dv}{dt} \quad (u \text{ & } v \text{ are vector function})$$

$$4. \frac{d(u \cdot v)}{dt} = u \cdot \frac{dv}{dt} + v \cdot \frac{du}{dt} \quad (u \text{ & } v \text{ are vector function})$$

$$5. \frac{d(uxv)}{dt} = u \times \frac{dv}{dt} + v \times \frac{du}{dt} \quad (u \text{ & } v \text{ are vector function})$$

### Chain rule

If  $r(t) = f(t)\mathbf{i} + g(t)\mathbf{j} + h(t)\mathbf{k}$  is a function of S then

$$\frac{dr}{ds} = \frac{dr}{dt} \cdot \frac{dt}{ds}$$

**Note:**  $u(t)$  is a function vector has constant length then

$$\bar{u} \cdot \overline{\frac{du}{dt}} = 0 \quad \text{or} \quad \bar{u} \perp \overline{\frac{du}{dt}}$$

Example: show that  $u(t) = \sin t \mathbf{i} + \cos t \mathbf{j} +$

*5k has constant length and is orthogonal to its derivative*

Solution//

$$\bar{u} \cdot \overline{\frac{du}{dt}} = 0$$

$$u(t) = \sin t \mathbf{i} + \cos t \mathbf{j} + 5\mathbf{k}$$

$$\frac{du}{dt} = \cos t \mathbf{i} - \sin t \mathbf{j} + 0\mathbf{k}$$

$$\bar{u} \cdot \overline{\frac{du}{dt}} = \sin t \cos t - \sin t \cos t + 0 = 0$$