

Assist.Prof. Dr. Ameer Mezher Hadi

Lecture. 2

Second Stage

Evaluation Lecturer



Al-Mustaqbal University

College of Health and Medical Techniques

Department of Medical Laboratory  
Technique

## *ESCHERICHIA COLI*

### Enteric Gram-negative rods: *Escherichia coli*

#### - The general characteristics of Enterobacteriaceae

- **Gram-negative** bacilli.
- Found as commensals in the intestinal tract of mammals.
- They are also referred to as **coliforms** or **enteric** bacteria.
- **Aerobic** and **facultative anaerobic** growth.
- Optimal growth normally at **37 C°**.
- Grow readily on **simple** media.
- **Ferment** wide range of **carbohydrates**.
- **According to the Lactose fermentation they are classified into:**
  - ❖ **Lactose fermenter** Fermentation of lactose to produce pink colonies on MacConkey's agar is characteristic of *Escherichia*, *Enterobacter* and *Klebsiella*.
  - ❖ **Non-lactose fermenter** *Salmonella*, *Shigella*, *Serratia*, *Proteus* and *Yersinia* do not ferment lactose and form pale colonies on MacConkey's agar.
  - ❖ **Late Lactose fermenters**, *Shigella sonnei*.
- **Oxidase-negative**.
- Some are **motile** (Motile except *Shigella* and *Klebsiella*).
- **Bile** tolerant and grow readily on **bile-salt** containing media, e.g. **MacConkey's** agar.
- **Some** of them produce **urease**. (which splits urea with release of ammonia).
- **Some** of them produce **Hydrogene sulphide**.

Assist.Prof. Dr. Ameer Mezher Hadi

Lecture. 2

Second Stage

Evaluation Lecturer




Al-Mustaqbal University

College of Health and Medical Techniques

Department of Medical Laboratory  
Technique

## ESCHERICHIA COLI

- Some of them **decarboxylase amino-acids**.
- Some of them **derive the indole ring** from the amino acid tryptophan
- **None-spore forming**.
- **None acid fast**.
- **Ferment** glucose with **acid** production
- **Reduce nitrates** into **nitrites**
- Non-capsulated except *Klebsiella*
- Non-fastidious



Enterobacteriaceae possess a variety of **antigens**:

- **lipopolysaccharide** somatic antigen('O'),
- **flagellar** antigen('H')
- **capsular polysaccharide** ('K') antigens.

### *Escherichia coli*

#### Morphology

- *E. coli* is Gram-negative (-ve) rod-shaped bacteria.
- It is 1-3 x 0.4-0.7 μm in size and 0.6 to 0.7 μm in volume.
- It is arranged singly or in pairs.
- It is motile due to **peritrichous** flagella (see classification of flagella ).
- Some **strains** are **non-motile**.
- **Some strains** may be **fimbriated**. The fimbriae are of type 1 (hemagglutinating & mannose-sensitive) and are present in both motile and non-motile strains.

Assist.Prof. Dr. Ameer Mezher Hadi

Lecture. 2

Second Stage

Evaluation Lecturer



Al-Mustaqbal University

College of Health and Medical Techniques

Department of Medical Laboratory  
Technique

## *ESCHERICHIA COLI*

- **Some strains** of *E. coli* isolated from extra-intestinal infections have a **polysaccharide capsule**.
- They are non-sporing.
- They have a thin cell wall with only 1 or 2 layers of peptidoglycan.
- They are facultative anaerobes.
- Growth occurs over a wide range of temperatures from 15-45°C.

### Antigenic Structure and Pathogenicity

- ✓ Specific **fimbriae (adhesins)** facilitate **adherence** to **mucosal surfaces** and **colonization** of the **intestinal** and **urinary** tracts.
- ✓ *E. coli* possesses 4 antigens; H, O, K and F.

#### A. Flagellar or (H) Antigen

- Heat and alcohol labile **protein**
- Present on the flagella
- Genus specific
- Present as monophasic
- 75 'H' antigens have been recognized

#### B. Somatic or (O) Antigen

- **Heat stable**, resistant to **boiling**.
- Occur on the surface of the outer membrane
- An integral part of the cell wall
- 173 'O' antigens have been recognized

#### C. Capsular or (K) Antigen

Assist.Prof. Dr. Ameer Mezher Hadi

Lecture. 2

Second Stage

Evaluation Lecturer



Al-Mustaqbal University

College of Health and Medical Techniques

Department of Medical Laboratory  
Technique

## *ESCHERICHIA COLI*

- Heat labile
- Acidic polysaccharide (polysaccharides that contain carboxyl groups, phosphate groups and/or sulfuric ester groups) antigen presents in the envelope
- Boiling removes the K antigen
- Inhibit phagocytosis
- 103 'K' antigens have been recognized

### **D. Fimbrial or (F ) Antigen**

- Heat labile proteins
- Present in the fimbriae
- K88, K99 antigens
- ✓ The **heat stable lipopolysaccharide (endotoxin)** "in the cell wall is liberated when Gram-negative bacteria lyse", resulting in production of **inflammatory mediators** and **complement activation** "plasma proteins that can be activated directly by pathogens or indirectly by pathogen-bound antibody". This results in **endotoxic shock** and **intravascular coagulopathy**.
- **Different protein toxins (exotoxins)** produced by *E. coli*.
- **Verocytotoxin-producing E. coli (VTEC), also known as (Shiga toxin-producing E. coli )(STEC)** particularly the **O157:H7** serotype, are an **important** cause of **diarrhoea** and **hemolytic uremic syndrome (HUS)**.
- **Enteropathogenic (EPEC):** cause of **infantile diarrhoea, non-invasive**.
- **Enterotoxigenic (ETEC):** **travelers' diarrhoea, non-invasive**.
- **Enteroinvasive (EIEC):** causes **dysentery-like illness**.

Assist.Prof. Dr. Ameer Mezher Hadi

Lecture. 2

Second Stage

Evaluation Lecturer



Al-Mustaqbal University

College of Health and Medical Techniques

Department of Medical Laboratory  
Technique

## *ESCHERICHIA COLI*

- **Enteroaggregative (EAEC): watery diarrhoea without fever.**

### **Pathogenicity of *E. coli***

**Most** infections (with the **exception of neonatal meningitis and gastroenteritis**) are **endogenous**; that is, the *E. coli* that are part of the patient's normal microbial flora are able to establish infection when the patient's defenses are compromised (e.g., through trauma or immune suppression).

This organism is associated with a variety of diseases, including **gastroenteritis** and **extra-intestinal infections** such as **UTIs, meningitis, and sepsis**.

### **Clinical Feature of *E. coli***

- 1. Gastroenteritis**
- 2. Urinary Tract Infection**
- 3. Sepsis**
- 4. Meningitis**

Assist.Prof. Dr. Ameer Mezher Hadi

Lecture. 2

Second Stage

Evaluation Lecturer




Al-Mustaqbal University


College of Health and Medical Techniques

Department of Medical Laboratory  
Technique

## *ESCHERICHIA COLI*



### Laboratory Diagnosis



*E. coli* on Blood Agar

Assist.Prof. Dr. Ameer Mezher Hadi

Lecture. 2

Second Stage

Evaluation Lecturer



Al-Mustaqbal University

College of Health and Medical Techniques

Department of Medical Laboratory  
Technique

## *ESCHERICHIA COLI*

---

1. Colonies are big, circular, gray, and moist.
2. Non-hemolytic colonies (gamma-hemolysis) (Above Figure) OR Beta ( $\beta$ )hemolytic (Below Figure) colonies are formed.
3. Many pathogenic strains are haemolytic on blood agar.



### *E. coli* on MacConkey Agar

1. Colonies are **circular, moist, smooth**, and of **entire** margin.
2. Colonies appear **flat** and **pink**.
3. They are **lactose** fermenting colonies.

Assist.Prof. Dr. Ameer Mezher Hadi

Lecture. 2

Second Stage

Evaluation Lecturer



Al-Mustaqbal University

College of Health and Medical Techniques

Department of Medical Laboratory  
Technique

## *ESCHERICHIA COLI*




*E. coli* on Eosin Methylene Blue (EMB) Agar

1. **Green Metallic** sheen colonies are formed.

### *E. coli* Biochemical Characters,

- **Glucose, Lactose, Mannitol, Maltose** fermented with **Acid and Gas**.



- Indole (+ve)
- Methyl Red (+ve)
- Voges Proskauer (-ve)
- Citrate (-ve)
- Urease not produced.
- H<sub>2</sub>S (-ve)
- Motility test (+ve)

Assist.Prof. Dr. Ameer Mezher Hadi

Lecture. 2

Second Stage

Evaluation Lecturer



Al-Mustaqbal University

College of Health and Medical Techniques

Department of Medical Laboratory  
Technique

## *ESCHERICHIA COLI*

---

**IMViC (+ + - -)**  
*E. coli*



Indole  
Positive

MR  
Positive

VP  
Negative

Citrate  
Negative