

Pediatric Anesthesia

By: Dr. Miaad Adnan

Dr. Basim Mohammed Jabbar.

2025

Introduction

It is often said that **paediatric** patients are '**not simply small adults**'. The truth is that from the premature neonate to the near-adult adolescent, children are very diverse.

Pediatric patients *involves* the following age groups :-

1. **Neonates** (0–1 months).
2. **infants** (1–12 months).
3. **toddlers** (12–24 months).
4. **young children** (2–12 years of age).

Pediatric anesthesia have differing anesthetic requirements. **physiological**, **anatomical**, and **pharmacological** characteristics of each group .

Indeed **infants** are at **much greater risk** of anesthetic morbidity and mortality than older children; **risk** is generally **inversely** proportional to **age**.

Estimation of weight

It is essential that every child is **weighed** prior to anaesthesia. This allows correct calculation of **drug doses** and selection of anaesthetic **equipment**. Weight can also be estimated from the age of the child from **standard growth charts**, from the **length of the child**, or using this **formula**:

| Age of child | Formula to estimate weight in kg |
|--------------|---|
| 0-12 months | $(0.5 \times \text{age in months}) + 4$ |
| 1-5 years | $(2 \times \text{age in years}) + 8$ |
| 6-12 years | $(3 \times \text{age in years}) + 7$ |



Anatomical & Physiological Differences

1-Respiratory System Differences

- The major anatomical differences affecting **airway management** in **neonates** and **infants** are:
 - **Relatively large head and prominent occiput**
 - **Small mandible**
 - **Relatively large tongue**
 - **Short neck**
 - **narrower nasal passages, an**
 - **anterior and cephalad larynx,**
 - **a longer epiglottis, and a shorter trachea.**
 - **Soft tracheal cartilages, easily compressed.**

These differences predispose to **airway obstruction**, particularly if the child's head is placed on a pillow, or the soft tissues on the floor of the mouth are compressed, or the head is hyperextended. **Ideally**, maintain the child's head in a **neutral** position, or **slightly** extended.

Anatomical & Physiological Differences

Anatomical differences affecting the larynx include:

- ❖ A **high**, anterior position of the larynx (level of **C3-4** in **infants** compared to **C5-6** in **adults**)
- ❖ A long, **U-shaped epiglottis**
- ❖ The **narrowest part** of the airway is at the **cricoid cartilage** (below the vocal cords).
- The **narrowest part** of the airway in **adults** is at the **vocal cords**.




FIGURE 42-1 Sagittal section of the adult (A) and infant (B) airway. (Reproduced with permission from Snell RS, Katz J. *Clinical Anatomy for Anesthesiologists*. New York, NY: Appleton & Lange; 1988.)

Anatomical & Physiological Differences

The major physiological differences in respiratory system.

- ❖ **Faster** respiratory rate.
- ❖ **Lower** lung compliance.
- ❖ **Greater** chest wall compliance.
- ❖ **Lower** functional residual capacity.
- ❖ **high** metabolic rate and oxygen consumption.

Note: children **tidal volume** is relatively **fixed** (5-7 ml.kg⁻¹), and the infant can **only** increase minute ventilation by increasing **respiratory rate**.

- ❖ **Apnoeas** are particularly **common** in **premature** and **ex-premature infants**, so monitor all babies for apneae after surgery; until they are 60 weeks post conception.
- ❖ If a mechanical ventilator is used, select the appropriate tidal volume and respiratory rate for age – **pressure control ventilation is preferred**.

Anatomical & Physiological Differences

2-Cardiovascular considerations

- ❖ Residual fetal circulation.
- ❖ Noncompliant left ventricle : so increase in **cardiac output** is achieved through an increase in **heart rate** (Heart-rate-dependent cardiac output).
- ❖ **Faster** heart rate: It is important to **avoid bradycardia**. This should be treated rapidly if it occur; the most **common cause** is **hypoxia**.
- ❖ **Lower** blood pressure

Age-related changes in vital signs.

| Age | Respiratory Rate | Heart Rate | Arterial Blood Pressure | |
|-----------|------------------|------------|-------------------------|-----------|
| | | | Systolic | Diastolic |
| Neonate | 40 | 140 | 65 | 40 |
| 12 months | 30 | 120 | 95 | 65 |
| 3 years | 25 | 100 | 100 | 70 |
| 12 years | 20 | 80 | 110 | 60 |

Anatomical & Physiological Differences

2-Cardiovascular considerations

- ❖ Activation of the **parasympathetic** nervous system by anesthetic **overdose**, or **hypoxia** can quickly trigger **bradycardia** and profound reductions in **cardiac output**.
- ❖ **Bradycardia** that can lead to **hypotension**, **asystole**, and intraoperative **death**.
- ❖ The **immature heart** is more sensitive to depression by **volatile anesthetics** and to **opioid-induced** bradycardia.
- ❖ **The main causes of neonatal bradycardia and cardiac arrest during anesthesia are:-**
 - ✓ **Respiratory causes** :- airway obstruction, bronchospasm, inadequate O₂ delivery.
 - ✓ **Pharmacology causes** :-inhalation agents, succinylcholine, anticholinesterase.
 - ✓ **Metabolic causes**:-hypothermia, anemia, hypoglycemia.
 - ✓ **Children are more** susceptible than adults to **cardiac arrhythmias**, **hyperkalemia**, **masseter spasm**, and **malignant hyperthermia associated with succinylcholine**.
 - ✓ When a child experiences cardiac arrest following administration of **succinylcholine**, immediate treatment for **hyperkalemia** should be instituted

Anatomical & Physiological Differences

3 . Metabolism & Temperature Regulation Differences.

Neonates promote greater **heat loss to the environment** and liable to hypothermia **Because:-**

1. **Thin** skin.
2. **Low** fat content.
3. **Greater** surface area relative to weight.
4. **Inadequately** warmed operating rooms (cold theater).
5. **Prolonged** wound exposure.
6. Administration of room temperature intravenous or irrigation fluid.
7. **Dry** anesthetic gases
8. Effects of anesthetic agents on temperature regulation center.

Even mild degrees of hypothermia can cause perioperative problems which including:-

- Delayed awakening from anesthesia.
- Cardiac irritability and arrest.
- Respiratory depression.
- Altered responses to anesthetics and Neuromuscular blockers, and other agents.

Anatomical & Physiological Differences

4-Renal & Gastrointestinal Function Differences

- The **total body water** is about **80%** of body weight at birth, gradually **decreasing** with **age**. fluid loss are more critical problem to them.
- **Immature kidney function** increases the importance of meticulous attention to fluid administration in the early days of life
- Neonates also have a relatively **increased** incidence of gastroesophageal **reflux**.
- The **immature liver** conjugates drugs and other molecules **less** readily early in life.

5-Glucose Homeostasis Differences

- Neonates have relatively **reduced** glycogen stores, predisposing them to **hypoglycemia**.

6-Pharmacological Differences: The main difference prolongs the clinical duration of action of drugs such as **thiopental** and **fentanyl**. this Bec:-

- 1.larger** pediatric intravascular and extracellular fluid **compartments** compare with adult . Neonates and infants have a proportionately greater total water content than adults (50–60%).
- 2.Immaturity** of hepatic **biotransformation** pathways,
- 3.Decreased protein** for drug binding .
- 4.Smaller muscle mass** in neonates **prolongs** or delaying redistribution of some drugs such as thiopental and fentanyl.

Some Pharmacological considerations

Volatile anesthetics

- Neonates are more sensitive to volatile agents than older children
- The minimum alveolar concentration (**MAC**) values are **decreased** in **neonates** but **increased** in infants and **children** compared to adults.

Sedatives and hypnotics

Children are particularly sensitive to sedative and hypnotic drugs such as **barbiturates** and **benzodiazepines** due to the **Immature hepatic biotransformation** and **Decreased protein binding** so these drugs should be used with caution, in weight appropriate doses, titrated according to effect.

Muscle relaxants

Neonates and **infants** are **more sensitive** to **non-depolarizing neuromuscular blocking** drugs **because** Immature neuromuscular junction.

A normal **loading dose** is given but **subsequent** doses should be **reduced**

ANAESTHETIC MANAGEMENT

Preoperative Preparation

- All children should be visited preoperatively by the anesthetist responsible for caring for them in the perioperative period.
- There is an increased incidence of airway problems during anaesthesia
- children are more at risk of laryngeal spasm, breath-holding and bronchospasm
- in the postoperative period the chance of post-intubation croup is increased.
- It is ***extremely important*** that the child is weighed before arrival in theatre, because body weight is the simplest and most reliable guide to drug dosage.
- Veins suitable for insertion of a cannula should be identified.

ANAESTHETIC MANAGEMENT

Preoperative Fasting

- Morbidity and mortality caused by aspiration of gastric contents are extremely rare in children undergoing elective surgery.
- **Prolonged** periods of **starvation** in children, especially the very young infant, are harmful.
- These children, who have a rapid turnover of fluids and a high metabolic rate, are at risk of developing **hypoglycemia** and **hypovolemia**.
- **Solids** (including breast and formula milk) should not be given for **at least 6 h before** the anticipated start of induction.
- In the **emergency** setting, e.g. the child who has sustained trauma shortly after ingesting food, **it is probably best (if possible) to wait 4 h before inducing anaesthesia**. Clearly, in this situation risk–benefit judgements have to be made.
- If it is surgically possible to wait 4 h, an i.v. infusion of a **glucose-containing solution** such as **5% dextrose** with **0.9% NaCl**, must be commenced and, if necessary, appropriate fluid resuscitation undertaken.

ANAESTHETIC MANAGEMENT

Intravenous Induction

- The same induction sequence can be used as in adults: a rapid-acting barbiturate (eg, thiopental, 3 mg/kg in neonates, 5–6 mg/kg in infants and children) or propofol (2–3 mg/kg) followed by a non-depolarizing muscle relaxant (eg, rocuronium, cisatracurium, atracurium, mivacurium, or succinylcholine).
- Atropine should be given intravenously prior to succinylcholine.
- It is important that children are accompanied into the anaesthetic room by someone with whom they are familiar.
- The appropriate monitoring should be applied as soon as possible **after** the start of anaesthesia.

ANAESTHETIC MANAGEMENT

- When inhalational induction is planned, clear, scented plastic masks are much more acceptable to small children than the traditional **Rendell–Baker rubber masks**.
- Clear masks allow respiration and the presence of vomit to be observed.
- An alternative to using a mask is cupping the hands over the face of the child while holding the T-piece, It is important to ensure that the flow of fresh gas is **directed away** from the child's **eyes** because anaesthetic gases may be irritant.
- When using a face mask, it is important that the soft tissue behind the chin is not pushed backwards by the fingers, thereby obstructing the airway. **The anaesthetist's fingers** should rest only on the **mandible**.



ANAESTHETIC MANAGEMENT

Airway Management

The Jackson–Rees modification of the Ayre's T-piece

- is the breathing system used traditionally for children under **20 kg** in weight.
- It has been designed to be **lightweight** with a **minimal** apparatus dead space.
- may be used for both **spontaneous** and **controlled** ventilation

The **open-ended** reservoir bag is used for manually controlled ventilation. This mode of ventilation is especially useful in the neonate and infant.

- Laryngeal mask airway (**LMA**) should be used **only** when it is planned that the child is to breathe **spontaneously** during surgery. It follows that **it is unwise** to use the device when neuromuscular blocking drugs are used.
- It is **mandatory** to intubate the trachea during **artificial** ventilation.
- Neonates with a tracheal tube must undergo artificial ventilation **in order to** *reduce the work of breathing*.
- Infants have a head which is **large** and a neck which is **short** relative to the size of the body. Instead of placing a pillow under the head, ***it is usually necessary to place a small pad or pillow under the torso.***

Airway Management

Tracheal intubation

For children over 1 year:

Appropriate tube internal diameter (ID) can be approximately estimated by the **formula**: $\text{age} / 4 + 4$.

Appropriate tube length in cm. can be approximately estimated by the **formula**: $\text{age} / 2 + 12$ oral (+15 for nasal).

In infants:

Appropriate tube ID sizes for preterm: <1500 g = 2.5 mm, 1500-3000 g = 3 mm, over 3000 g = 3.5 mm.

Oral length in cm. is given by the **formula** ($6 + \text{weight in kg}$).

Laryngeal mask airway (LMA): They are useful in short procedures with spontaneous ventilation. They have less resistance than endotracheal tubes and are of considerable use for insertion of fiberoptic bronchoscopes. Approximate sizes are:

- 1 for less than 6.5 kg.
- 2 for 6.5-20 kg.
- 2.5 for 20-30 kg.
- 3 for 30 kg and above.

A size 1.5 is also available.

The armored versions have reduced risk of kinking and are longer and narrower.

Airway Management

If too **large** a tube is selected, the tracheal mucosa is **damaged** and the child may develop **post-intubation croup**;

If it is **too small**, excessive **leak** makes effective positive pressure ventilation impossible

Generally, **cuffed tubes** are used only in children **above** the age of 8 years.

Adequate **preoxygenation** may help prevent hypoxemia.

A **prominent occiput** tends to place the head in a *flexed position* prior to intubation. This is easily corrected by *slightly elevating the shoulders with towels*.

Straight laryngoscope **blades** aid intubation of the **anterior larynx** in neonates, infants, and young children.

Mucosal trauma from trying to force a tube through the cricoid cartilage can cause postoperative **edema, stridor, croup, and airway obstruction**.

Inhalational Induction

- Modern **potent volatile** anesthetics can render small children unconscious within minutes. This is usually easier in children who have been **sedated** prior to entering the operating room and who are sleepy enough to be anesthetized without ever knowing what has happened (***steal induction***).
- ***Inhalational Induction***
- Equipment appropriate for age and size should be selected.
- Typically, the child is coaxed into breathing an odorless mixture of nitrous oxide (70%) and oxygen (30%). **Sevoflurane** is added to the anesthetic gas mixture in **0.5%** increments every **three** to **five** breaths.
- **Sevoflurane** consider ***the agent of choice for inhalation induction***.

Maintenance

- Ventilation is usually **controlled** during anesthesia of neonates and infants.
- During **spontaneous** ventilation, even the low resistance of a circle system can become a significant obstacle for a sick neonate to overcome.
- For patients weighing **less than 10 kg, it's** preferred to use the **Mapleson D** circuit or the **Bain system** because of their low resistance and light weight.
- **The circle system** can be safely used in patients of all ages if ventilation is controlled.
- Anesthesia can be **maintained** in pediatric patients with the same agents as in adults.

Maintenance

- **Isoflurane** Can be use following a **sevoflurane** induction to help reduce the likelihood of postoperative delirium or agitation on emergence.
- If **sevoflurane** is continued for maintenance, administration of an **opioid** (eg, fentanyl 1–1.5 mg/kg) 15–20 min before the end of the procedure can reduce the incidence of emergence delirium and agitation.
- Although the **MAC is higher in children** than in adults, neonates may be particularly susceptible to the cardiodepressant effects of general anesthetics. Nondepolarizing muscle relaxants are often required for optimal surgical conditions.

Perioperative Fluid Requirements

- Fluid therapy can be divided into **maintenance**, **deficit**, and **replacement** requirements.

MAINTENANCE FLUID REQUIREMENTS

Maintenance requirements for pediatric patients can be determined by the **4:2:1 rule**: 4 mL/kg/h for the first 10 kg of weight, 2 mL/kg/h for the second 10 kg, and 1 mL/kg/h for each remaining kilogram.

DEFICITS

- In addition to a maintenance infusion, any preoperative fluid deficits must be replaced. Calculated as (**maintenance fluid x starvation hours**) For example, if a **5-kg** infant has not received oral or intravenous fluids for **4 h** prior to surgery, a deficit of 80 mL has accrued ($5 \text{ kg} \times 4 \text{ mL/kg/h} \times 4 \text{ h}$).
- Preoperative fluid deficits are typically administered with hourly maintenance requirements in aliquots of **50%** in the **first hour** and **25%** in the **second** and **third** hours. In the example above, a total of 60 mL would be given in the first hour ($80/2 + 20$) and **40** mL in the second and third hours ($80/4 + 20$).
- **Preoperative fluid deficits are usually replaced with a balanced salt solution (eg, lactated Ringer's injection) or 1/2 normal saline.**

REPLACEMENT REQUIREMENTS

- Replacement can be subdivided into blood loss and third-space loss.

Blood Loss

- **Blood** loss is typically replaced with non-glucose-containing crystalloid (e.g., 3 mL of lactated Ringer's injection for each milliliter of blood lost) or colloid solutions (eg, 1 mL of 5% albumin for each milliliter of blood lost).

THANK YOU