

# Paranasal Sinuses

Dr Zaid Saad Al Nasrawi

Dr Lamees Abd AlRaheem

# Anatomy


**The paranasal sinuses** are air-filled cavities that open into the nasal cavity, mostly into the middle meatus of the nose.

**The maxillary sinuses** occupy the cheeks.

**The ethmoid labyrinth** consists of a number of air cells lying between the orbit and the lateral wall of the nose.

**The frontal sinus** is an ethmoid air cell that has migrated into the frontal bone, and it is connected to the middle meatus of the nose via the frontonasal duct.

**The sphenoid sinus** is posterior to the ethmoid labyrinth, inferior to the pituitary fossa.




Frontal sinus

Sphenoid sinus

Ethmoid air cells

Maxillary sinus

Lateral







Frontal sinus

Sphenoid sinus

Ethmoid air cells

Maxillary sinus





# Diseases of the paranasal sinuses


- Any sinus may become **infected**.
- The **most commonly involved** is the **maxillary sinus**.
- **Pain** arising from the **maxillary sinus** is felt in the cheek, that from the **ethmoid labyrinth** is felt over the nasal bridge, and frontal sinus pain is felt in the forehead. **Sphenoid sinus** pain is said to be maximal at the **vertex**.
- **Acute sinusitis** is most commonly caused by **Strep. pneumoniae** or **H. influenzae**, and typically follows an upper respiratory infection. **Gram negative organisms** may cause sinusitis related to a dental abscess. In some parts of the world, **fungal infection** is not uncommon.

Acute sinusitis is usually managed medically.

Chronic sinusitis may result from failure of resolution of acute infection or may arise insidiously.

Surgical treatment is frequently required and includes enlargement of the natural ostium of the maxillary sinus, often with clearance of infected ethmoid cells.

Frontal and sphenoid sinusitis are much less common. Infection may spread from the sinuses, usually the ethmoid or frontal sinuses, to involve other areas such as the cranial cavity or orbit .



# Complications:

1. Mucus retention cyst, polyps
2. Mucocele
3. Osteomyelitis
4. Cavernous sinus thrombosis
5. Intracranial extension ( Empyema, Cerebritis , Abscess )
6. Orbital complications

# Tumors

The most common malignant neoplasm found in the paranasal sinuses is squamous carcinoma.

Adenocarcinomas are seen in workers in the furniture industry.

The most common sites of origin are the maxillary and ethmoid sinuses.

Unfortunately, the disease has often spread beyond the primary site at presentation.

These relatively uncommon tumors are managed by a combination of surgery and radiotherapy, or by local surgery and topical chemotherapy.

# Imaging of the paranasal sinuses

Evaluation of the paranasal sinuses is often performed in a purely clinical fashion, without the need for imaging.

However, in certain instances imaging may be deemed valuable or even necessary in helping to solve a diagnostic dilemma, confirm a suspected diagnosis, evaluate the extent of a known condition, or assess for an underlying cause of the condition.

X-ray , Computed tomography (CT) and magnetic resonance imaging (MRI) can be useful in confirming a suspected diagnosis or providing additional information regarding causes or complications.


CT and MRI play complementary roles in evaluating the rare tumors that may involve the paranasal sinuses.

# Plain radiograph ( x-ray )

Basic radiographic features are **occipitofrontal view** (Caldwell view ) , **occipitomental ( OM )** or **Waters view** and **lateral view**

Sinuses are normally filled with air, so the passages will appear black on an X-ray of healthy sinuses.

**A gray or white area** on an X-ray of the sinuses indicates a problem. This is most often due to inflammation or a buildup of fluid in the sinuses.






# CT scan

The CT paranasal sinus protocol serves as an examination for the assessment of the study of the mucosa and bone system of the sinonasal cavities.

It is usually performed as a non-contrast study.

**Typical indications include the following:**

1. Inflammatory disease ( acute rhinosinusitis , gas-fluid levels , mucosal disease , chronic sinusitis , cysts and polyps , mucoceles )
2. Foreign body
3. Malignancy
4. Preoperative assessment



In neoplastic, inflammatory and infectious disorders, a CT scan of the paranasal sinuses is performed to demonstrate bony erosions, osteolytic lesions, and calcifications.

If neoplasia is suspected, the use of an intravenous contrast medium is indicated.

CT is of value for determining anatomic landmarks and variants. This information is of vital importance to the ENT-surgeon.

In addition, we need it to identify erosive processes and acquired developmental deficiencies of the bone.

CT is also excellent for determining whether there is intraorbital extension of sino-nasal disease in the ventral 2/3 of the orbit.

If additional imaging is necessary, orbital MRI is the next step. The real value of unenhanced CT is the following: if you see an opacified sinus with hyperdense contents, it is usually a sign of benign disease. Tumor is not hyper-dense.

The hyperdensity is due to one or a combination of the following:

1. inspissated secretions
2. fungus
3. blood




# MRI

MRI is extremely helpful in complicated sinonasal disease.

MRI can discern secretions and mucosa from masses.

When you understand the signal characteristics, you are readily able to distinguish soft tissue masses from inspissated secretions.

The signal intensity of secretions can vary and depends mainly on the water-to-protein ratio and viscosity. Different protein contents result in different signal intensities on T1 and T2W images.



Fungus usually has a high protein content of more than 28% and can mimic an aerated sinus because it is low on T1- and T2WI.

MRI is also useful for determining invasion of the skull base.

Involvement of the skull base is seen as replacement of the high signal of the fatty marrow on T1WI by hypointense signal of the tumor.

Also look for foraminal extension, whether by perineural spread or direct invasion of the tumor

MRI is also the study of choice for detecting intracranial extension of sinonasal disease.

In general bright signal on T2 is a sign of benign disease, since fluid and mucosal disease usually have a high water content. Secretions do not have solid enhancement.

If you have an enhancing mass, you must rule out tumor.

After the administration of i.v. contrast there is only enhancement of the circumferential mucosa and no solid enhancement.

In complicated cases both CT and MR are needed to demonstrate the extension of the disease.