



Al-Mustaqbal University / College of Engineering & Technology  
Department of Techniques of Fuel and Energy Engineering

Class:one

Subject: Analytical chemistry / Code: UOMU0206011

Lecturer: M.Sc. Zahraa ALjassar

1<sup>st</sup>term – Lecture 4 (pH for the acidic solutions)



## pH for the acidic solution

In order to find the numeric value of the level of acidity or basicity of a substance, the pH scale (where in pH stands for ‘potential of hydrogen’) can be used. The pH scale is the most common and trusted way to measure how acidic or basic a substance. A pH scale measure can vary from 0 to 14, where 0 is the most acidic and 14 is the most basic a substance can be. Another way to check if a substance is acidic or basic is to use litmus paper. There are two types of litmus paper available that can be used to identify acids and bases – red litmus paper and blue litmus paper. Blue litmus paper turns red under acidic conditions and red litmus paper turns blue under basic or alkaline conditions.



Al-Mustaqbal University / College of Engineering & Technology  
Department of Techniques of Fuel and Energy Engineering

Class:one

Subject: Analytical chemistry / Code: UOMU0206011

Lecturer: M.Sc. Zahraa ALjassar

1<sup>st</sup>term – Lecture 4 (pH for the acidic solutions)



## pH and pOH

pH scale is a commonly used scale to measure the acidity or the basicity of a substance. The possible values on the pH scale range from 0 to 14. Acidic substances have pH values ranging from 1 to 7 (1 being the most acidic point on the pH scale) and alkaline or basic substances have pH values ranging from 7 to 14.

A perfectly neutral substance would have a pH of exactly 7.

pH which is an abbreviation of ‘potential for hydrogen’ or ‘power of hydrogen’ of a substance can be expressed as the negative logarithm (with base 10) of the hydrogen ion concentration in that substance.

Similarly,

the pOH of a substance is the negative logarithm of the hydroxide ion concentration in the substance.



Al-Mustaqbal University / College of Engineering & Technology  
Department of Techniques of Fuel and Energy Engineering

Class:one

Subject: Analytical chemistry / Code: UOMU0206011

Lecturer: M.Sc. Zahraa ALjassar

1<sup>st</sup>term – Lecture 4 (pH for the acidic solutions)



**These quantities can be expressed via the following formulae:**

$$\text{pH} = -\log [H^+]$$

$$\text{pH} = -\log [OH^-]$$

Or

$$\text{POH} = -\log [OH^-]$$

**Both pH and pOH are related to each other. pH is inversely proportional to pOH; pH increases with decreasing pOH.**

**Relation between p [H<sup>+</sup>]**

$$\text{PH} \propto \frac{1}{\text{POH}}$$

**and p [OH] :**

$$\text{pH} + \text{pOH} = 14$$



## pH – A Measure of Acidity

$$\text{pH} = -\log [\text{H}^+]$$

### Solution Is

### At 25°C

neutral	$[\text{H}^+] = [\text{OH}^-]$	$[\text{H}^+] = 1.0 \times 10^{-7}$	pH = 7
acidic	$[\text{H}^+] > [\text{OH}^-]$	$[\text{H}^+] > 1.0 \times 10^{-7}$	pH < 7
basic	$[\text{H}^+] < [\text{OH}^-]$	$[\text{H}^+] < 1.0 \times 10^{-7}$	pH > 7





## Other important relationships

$$\text{pOH} = -\log [\text{OH}^-]$$

$$[\text{H}^+][\text{OH}^-] = K_w = 1.0 \times 10^{-14}$$

$$-\log [\text{H}^+] - \log [\text{OH}^-] = 14.00$$

$$\text{pH} + \text{pOH} = 14.00$$

### The pHs of Some Common Fluids

Sample	pH Value
Gastric juice in the stomach	1.0–2.0
Lemon juice	2.4
Vinegar	3.0
Grapefruit juice	3.2
Orange juice	3.5
Urine	4.8–7.5
Water exposed to air*	5.5
Saliva	6.4–6.9
Milk	6.5
Pure water	7.0
Blood	7.35–7.45
Tears	7.4
Milk of magnesia	10.6
Household ammonia	11.5



pH Meter



### Example 5:

The concentration of  $H^+$  ions in a solution is  $3.2 \times 10^{-4} M$ . The solution was left for a while, it was found that the hydrogen ion concentration equal to  $1.0 \times 10^{-3} M$ . Calculate the pH of the solution on these two occasions.

$$pH = -\log [H^+] \quad [H^+] = 3.2 \times 10^{-4} M$$

$$pH = -\log [H^+]$$

$$= -\log (3.2 \times 10^{-4}) = 3.49$$

On the second occasion,  $[H^+] = 1.0 \times 10^{-3} M$ ,

so that

$$pH = -\log (1.0 \times 10^{-3}) = 3.00$$



### Example 7:

In a NaOH solution  $[\text{OH}^-]$  is  $2.9 \times 10^{-4} \text{ M}$ . Calculate the pH of the solution.

$$\begin{aligned}\text{pOH} &= -\log [\text{OH}^-] \\ &= -\log (2.9 \times 10^{-4}) \\ &= 3.54\end{aligned}$$

$$\begin{aligned}\text{pH} + \text{pOH} &= 14.00 \\ \text{pH} &= 14.00 - \text{pOH} \\ &= 14.00 - 3.54 = \mathbf{10.46}\end{aligned}$$

Alternatively, we can use  $K_w = [\text{H}^+][\text{OH}^-]$  to calculate  $[\text{H}^+]$ , and then we can calculate the pH from the  $[\text{H}^+]$ . *Try it.*

$$[\text{H}^+] = \frac{K_w}{[\text{OH}^-]} \longrightarrow \text{pH} = -\log [\text{H}^+]$$




## Strength of acids and bases





Strong Electrolyte – 100% dissociation



Weak Electrolyte – not completely dissociated






**Strong Acids** are strong electrolytes








## **Weak Acids** are weak electrolytes





## **Strong Bases** are strong electrolytes







## Strong Acid (HCl)





## Weak Acid (HF)





## Weak Bases are weak electrolytes



## Conjugate acid-base pairs:

- The conjugate base of a strong acid has no measurable strength.
- $\text{H}_3\text{O}^{+}$  is the strongest acid that can exist in aqueous solution.
- The  $\text{OH}^{-}$  ion is the strongest base that can exist in aqueous solution.



## Relative strength of acids and bases

	Acid	Base	
<div>Strongest acids</div> <div><div></div></div>	HClO <sub>4</sub>	ClO <sub>4</sub> <sup>-</sup>	<div>Weakest bases</div> <div><div></div></div>
	H <sub>2</sub> SO <sub>4</sub>	HSO <sub>4</sub> <sup>-</sup>	
	HI	I <sup>-</sup>	
	HBr	Br <sup>-</sup>	
	HCl	Cl <sup>-</sup>	
	HNO <sub>3</sub>	NO <sub>3</sub> <sup>-</sup>	
	H <sub>3</sub> O <sup>+</sup>	H <sub>2</sub> O	
	HSO <sub>4</sub> <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	
	H <sub>2</sub> SO <sub>3</sub>	HSO <sub>3</sub> <sup>-</sup>	
	H <sub>3</sub> PO <sub>4</sub>	H <sub>2</sub> PO <sub>4</sub> <sup>-</sup>	
	HNO <sub>2</sub>	NO <sub>2</sub> <sup>-</sup>	
	HF	F <sup>-</sup>	
	HC <sub>2</sub> H <sub>3</sub> O <sub>2</sub>	C <sub>2</sub> H <sub>3</sub> O <sub>2</sub> <sup>-</sup>	
	Al(H <sub>2</sub> O) <sub>6</sub> <sup>3+</sup>	Al(H <sub>2</sub> O) <sub>5</sub> OH <sup>2+</sup>	
	H <sub>2</sub> CO <sub>3</sub>	HCO <sub>3</sub> <sup>-</sup>	
	H <sub>2</sub> S	HS <sup>-</sup>	
	HClO	ClO <sup>-</sup>	
	HBrO	BrO <sup>-</sup>	
	NH <sub>4</sub> <sup>+</sup>	NH <sub>3</sub>	
	HCN	CN <sup>-</sup>	
HCO <sub>3</sub> <sup>-</sup>	CO <sub>3</sub> <sup>2-</sup>		
H <sub>2</sub> O <sub>2</sub>	HO <sub>2</sub> <sup>-</sup>		
HS <sup>-</sup>	S <sup>2-</sup>		
H <sub>2</sub> O	OH <sup>-</sup>	<div>Strongest bases</div>	




### Example 8

Calculate the pH of a :

(a)  $1.0 \times 10^{-3} \text{ M HCl}$  solution

(b)  $0.020 \text{ M Ba(OH)}_2$  solution




	HCl(aq)	$\rightarrow$	H <sup>+</sup> (aq)	+	Cl <sup>-</sup> (aq)
Initial (M):	$1.0 \times 10^{-3}$		0.0		0.0
Change (M):	$-1.0 \times 10^{-3}$		$+1.0 \times 10^{-3}$		$+1.0 \times 10^{-3}$
Final (M):	0.0		$1.0 \times 10^{-3}$		$1.0 \times 10^{-3}$

$$\begin{aligned} [\text{H}^+] &= 1.0 \times 10^{-3} \text{ M} \\ \text{pH} &= -\log (1.0 \times 10^{-3}) \\ &= 3.00 \end{aligned}$$



### (b) $\text{Ba}(\text{OH})_2$ is a strong base



$$[\text{OH}^-] = 0.040 \text{ M}$$

$$\text{pOH} = -\log 0.040 = 1.40$$

$$\begin{aligned} \text{pH} &= 14.00 - \text{pOH} \\ &= 14.00 - 1.40 \\ &= 12.60 \end{aligned}$$