

## Lecture (2)



**Example 8:** Find the line L1 passes through the point  $P(1,2)$  and parallel the line  $L2: x + 2y = 3$ .

**SOL:**

$$L1: P(1,2) \quad M=???$$

$$L2: x + 2y = 3.$$

$L1$  parallel the line  $L2$  so that  $m1=m2$ .

$$x + 2y = 3$$

$$y = -\frac{1}{2}x + \frac{3}{2}$$

then  $m2 = -\frac{1}{2}$  so that  $m1 = -\frac{1}{2}$

$$y = y_1 + m(x - x_1)$$

$$y = 2 + \left(-\frac{1}{2}\right)(x - 1)$$

$$y = 2 + \left(-\frac{1}{2}x + \frac{1}{2}\right)$$

$$y = -\frac{1}{2}x + \frac{5}{2}$$

**H.W:**

**Find the line L1 passes through the point  $(-2,2)$  and perpendicular to the line  $L2 : 2x + y = 4$ .**

## Lecture (2)

The Distance from a Point to a Line:

The distance (d) between the line L is  $Ax + By + C = 0$  and the point  $P(x, y)$ :

$$d = \frac{|Ax_1 + By_1 + C|}{\sqrt{A^2 + B^2}}$$

**Example 9:** Find the distance from the point  $P(2,1)$  to the line  $y = x + 2$

**SOL:**

1- put the line in the general form  $Ax + By + C = 0$

$$y = x + 2$$

$$-x + y - 2 = 0$$

so that  $A = -1$  ,  $B = 1$  ,  $C = -2$  ,  $x_1 = 2$  ,  $y_1 = 1$

$$d = \frac{|Ax_1 + By_1 + C|}{\sqrt{A^2 + B^2}} = \frac{|-1 * (2) + 1 * (1) + (-2)|}{\sqrt{(-1)^2 + (1)^2}}$$

$$= \frac{|-3|}{\sqrt{2}} = \frac{3}{\sqrt{2}}$$

**H.W:**

1-Find the distance from the point  $P(3,2)$  to the line  $y = 3x - 4$ .

2-Find the distance from the point  $P(-4,1)$  to the line  $y = -2x + 1$ .

3- Find the following:

- The slope of the line  $2x + 3y - 5 = 0$ ?
- The distance from the above line to the point  $P(-1,0)$ .




## الدوال Functions

### DEFINITION: Function

A **function** is a set D (domain) to a set R (range) is a rule that assigns to unique (single) element  $f(x) \in R$  to each element  $x \in D$ .

$F: X \rightarrow F(X)$  it means that f sends x to  $f(x)=y$



$D$  = domain set

$Y$  = set containing the range

- The set of x is called the "Domain" of the function (D<sub>f</sub>).
- The set of y is called the "Range" of the function (R<sub>f</sub>).

**Domain (D<sub>f</sub>)**: is the set of all possible inputs (x-values).

**Range (R<sub>f</sub>)**: is the set of all possible outputs (y-values).

**Note:** To find Domain (D<sub>f</sub>) and the Range (R<sub>f</sub>) the following points must be noticed:

- 1- The denominator in a function must not equal zero (never divide by zero).
- 2- The values under even roots must be positive.



**Examples:** Find the Domain (Df) and Range (Rf) of the following functions:

1-  $y = f(x) = \frac{1}{x}$

**Sol:** denominator must not equal zero

$$x \neq 0$$

✓  $Df = R \setminus \{0\}$

To find Rf : we must convert the function from  $y=f(x)$  into  $x=f(y)$ .

$$y = \frac{1}{x} \rightarrow x = \frac{1}{y}$$

✓  $Rf = R \setminus \{0\}$ .

2-  $y = \sqrt{3 - x}$

$$3 - x \geq 0 \rightarrow 3 \geq x$$

✓  $Df = \{x \in R \mid x \leq 3\}$

To find Rf : we must convert the function from  $y=f(x)$  into  $x=f(y)$ .

$$y = \sqrt{3 - x}$$

$$y^2 = 3 - x$$

$$x = 3 - y^2$$

✓  $Rf = \{y \in R\}$ .

**H.W:** Find the Domain (Df) and Range (Rf) of the following functions:

1-  $y = \frac{1}{x^2}$

2-  $y = 2x^2$

3-  $y = \sqrt{5 - 2x}$

## Lecture (2)

**Sums, Difference, Product and Quotients of Functions:****جمع، طرح، ضرب وقسمة الدوال**

**Definition:** If  $F$  and  $G$  are functions, then we define the functions

- ✓ Sum  $\rightarrow (F+G)(x) = F(x) + G(x)$
- ✓ Difference  $\rightarrow (F - G)(x) = F(x) - G(x)$
- ✓ Product  $\rightarrow (F * G)(x) = F(x) * G(x)$
- ✓ Quotient  $\rightarrow (F / G)(x) = F(x) / G(x)$ , where  $g(x) \neq 0$

**Example 1: Combining Functions Algebraically**

The function defined by the formulas

$$f(x) = \sqrt{x} \text{ and } g(x) = \sqrt{1-x}$$

Function	Formula
$f + g$	$(f + g)(x) = \sqrt{x} + \sqrt{1-x}$
$f - g$	$(f - g)(x) = \sqrt{x} - \sqrt{1-x}$
$g - f$	$(g - f)(x) = \sqrt{1-x} - \sqrt{x}$
$f \circ g$	$(f \circ g)(x) = f(x)g(x) = \sqrt{x(1-x)} = \sqrt{x-x^2}$
$f/g$	$\frac{f}{g}(x) = \frac{f(x)}{g(x)} = \sqrt{\frac{x}{1-x}}$
$g/f$	$\frac{g}{f}(x) = \frac{g(x)}{f(x)} = \sqrt{\frac{1-x}{x}}$

**H.W: Combining Functions Algebraically** The function defined by the formulas  $f(x) = 3x$  and  $g(x) = 1 - x^2$ .