




DNA Transcription

Lac5



Transcription


- Transcription, is the process of creating an equivalent RNA copy of a sequence of DNA.
- Transcription is the first step leading to **gene expression**.

Transcription is the first of several steps of DNA based gene expression in which a particular segment of DNA is copied into RNA by the enzyme RNA polymerase. Both DNA and RNA are nucleic acids

- Double stranded DNA must be **TRANSCRIBED** into Single stranded RNA
- **.1mRNA “messenger”**
- made from DNA in nucleus...travels out of nucleus and finds a ribosome.
- **.2tRNA “transfer”**
- brings amino acids to the ribosomes; found in cytoplasm
- **.3rRNA “ribosomal”**
- part of the ribosome; this is where proteins are made

How does the DNA get made into RNA and that made into Protein?


During transcription, the enzyme RNA polymerase (green) uses DNA as a template to produce a pre-mRNA transcript (pink). The pre-mRNA is processed to form a mature mRNA molecule that can be translated to build the protein molecule (polypeptide) encoded by the original gene



Transcription in prokaryote

- RNA molecules are produced by copying part of a nucleotide sequence of DNA into a complementary sequence in RNA. This Process is called transcription.
- Transcription requires the enzyme RNA polymerase.
- During transcription, a DNA sequence is read by RNA polymerase, which produces a complementary, antiparallel RNA strand.
- Transcription results in an RNA complement that includes uracil (U) instead of thymine (T).



Coding and Non-Coding Strands




Stages of Transcription in Prokaryote

- Initiation
- RNA polymerase attaches to the DNA molecule and moves along the DNA strand until it recognizes a promoter sequence (Sequence on DNA 10 and 35 bases upstream of start site). These are known as the transcription start sites. The DNA double helix then unwinds and all the bases on each of the DNA strands are exposed. This acts as a template for a new mRNA strand.
- Elongation
- Ribonucleotides are added to the template strand that enables the growth of mRNA.
- Termination
- RNA polymerase encounters a terminator sequence and the transcription stops. RNA polymerase then releases the DNA template.

Stages of Transcription





(5') CGCTATAGCGTTT (3')

DNA nontemplate (coding) strand

(3') GCGATATCGCAA (5')

DNA template strand

(5') CGCUAUAGCGUUU (3')

RNA transcript


Figure 26-2

Lehninger Principles of Biochemistry, Fifth Edition

© 2008 W.H. Freeman and Company


RNA polymerase in prokaryotic

- In most prokaryotes, a **single RNA polymerase species transcribes all types of RNA**. RNA polymerase "core" from E. coli consists of five subunits: two alpha (α) subunits of 36 kDa, a beta (β) subunit of 150 kDa, a beta prime subunit (β') of 155 kDa, and a small omega (ω) subunit.
- **Sigma (σ) factor**
 - *Smaller protein
 - *Guides RNA polymerase to target DNA sequence
 - *Sigma factor released after open complex (unwinds DNA at promoter)



Eukaryotic Transcription

- Promoters
- Much more complex than those found in bacteria.
- These are consensus sequences located at the upstream regions of Coding strand.
- Mutation of this region usually significantly lowers the rate of transcription.



Post-Transcriptional Modifications


Posttranscriptional modifications to eukaryotic pre-mRNA

| Modification | Function |
|--|--|
| Addition of 5' cap | Facilitates binding of ribosome to 5' end of mRNA, increases mRNA stability, enhances RNA splicing |
| 3' cleavage and addition of poly(A) tail | Increases stability of mRNA, facilitates binding of ribosome to mRNA |
| RNA splicing | Removes noncoding introns from pre-mRNA, facilitates export of mRNA to cytoplasm, allows for multiple proteins to be produced through alternative splicing |
| RNA editing | Alters nucleotide sequence of mRNA |

RNA polymerase in Eukaryote

| Name | Can be found in | Transcribes |
|--------------------|-----------------|---|
| RNA Polymerase I | All Eukaryotes | Large rRNAs |
| RNA Polymerase II | All Eukaryotes | mRNA, snoRNAs, some snRNAs and miRNAs |
| RNA Polymerase III | All Eukaryotes | tRNAs, small rRNAs, some snRNAs and miRNAs |
| RNA Polymerase IV | Plants | some siRNAs |
| RNA Polymerase V | Plants | RNAs important in heterochromatin formation |

طلبة



Prokaryotic vs Eukaryotic Transcription

| Prokaryotic Transcription | Eukaryotic Transcription |
|---|---|
| Transcription and translation occur simultaneously | Transcription and translation don't occur simultaneously. |
| Prokaryotic transcription occurs in the cytoplasm | Eukaryotic transcription occurs in the nucleus and translation occurs in the cytoplasm. |
| RNAs are released and processed in the cytoplasm | RNAs are released and processed in the nucleus |
| RNA polymerases are a complex of five polypeptides. | RNA polymerases are a complex of 10 -15 polypeptides. |



Thank you

