



جامعة المستقبل  
AL MUSTAQL UNIVERSITY

كلية العلوم  
قسم الذكاء الاصطناعي



Lecture (5)

MATRICES AND DETERMINANTS

المادة : الرياضيات

المرحلة : الاولى

اسم الاستاذ: م.م رياض ثائر احمد



**k7kgjyvz**

<https://classroom.google.com/c/ODM3NjM4NTg2MzY3?cjc=k7kgjyvz>



## Matrices and Determinants

A matrix is a rectangular array of elements (scalars) from a field. The order, or size, of a matrix is specified by the number of rows and the number of columns, i.e.  $A$  an “ $m$  by  $n$ ” matrix has  $m$  rows and  $n$  columns, and the element in the  $i$ th row and  $j$ th column is often denoted by  $a_{ij}$ :

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

A vector is a matrix with a single row (or column) of  $n$  elements, i.e. the column vector is:-

$$A = \begin{bmatrix} a_1 \\ a_2 \\ \cdot \\ \cdot \\ a_n \end{bmatrix} \quad \text{and row vector is} \quad A = [a_1 \ a_2 \ \dots \ a_n]$$

The matrix is square if the number of rows and columns are equal (i.e.  $m = n$ ) and the elements  $a_{ij}$  of a square matrix are called the main diagonal.

$$\text{The identity matrix: } I = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{bmatrix} \quad \text{is square matrix}$$

with one in each main diagonal position and zeros else.



The diagonal matrix  $D = \begin{bmatrix} a_1 & 0 & \dots & 0 \\ 0 & a_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_n \end{bmatrix}$  has the elements  $a_1, a_2, \dots, a_n$  in its main diagonal position and zeros in all other locations, some of the  $a_i$  may be zero but not all.

A  $n \times n$  triangular matrix has the pattern:-

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ 0 & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix}$$

lower triangular matrix

or

$$\begin{bmatrix} a_{11} & 0 & \dots & 0 \\ a_{21} & a_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

upper triangular matrix

The  $m \times n$  null matrix:-  $0 = \begin{bmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 \end{bmatrix}$  has zero in each of its positions.

## Elementary operations with matrices and vectors

1. Equality:- Two  $m \times n$  matrices and  $A$  and  $B$  are said to be equal if:  $a_{ij} = b_{ij} \quad \forall$  pairs of  $i$  and  $j$ .

EX-1 – Find the values of  $x, y$  for the following matrix equation:

$$\begin{bmatrix} x - 2y & 0 \\ -2 & 6 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ -2 & x + y \end{bmatrix}$$

Sol. –



$$\begin{array}{l} x - 2y = 3 \\ x - 2y = 3 \quad \dots \dots (1) \\ x + y = 6 \quad \dots \dots (2) * 2 \end{array} \Rightarrow \begin{array}{l} 2x + 2y = 12 \\ 3x = 15 \Rightarrow x = 5 \end{array}$$

$$\text{substitution } x = 5 \text{ in (2)} \Rightarrow 5 + y = 6 \Rightarrow y = 1$$

2. Addition:- The sum of two matrices of like dimensions is the matrix of the sum of the corresponding elements. If:-

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1n} \\ b_{21} & b_{22} & \dots & b_{2n} \\ \dots & \dots & \dots & \dots \\ b_{m1} & b_{m2} & \dots & b_{mn} \end{bmatrix}$$

then

$$A \mp B = \begin{bmatrix} a_{11} \mp b_{11} & a_{12} \mp b_{12} & \dots & a_{1n} \mp b_{1n} \\ a_{21} \mp b_{21} & a_{22} \mp b_{22} & \dots & a_{2n} \mp b_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} \mp b_{m1} & a_{m2} \mp b_{m2} & \dots & a_{mn} \mp b_{mn} \end{bmatrix}$$

thus:

- 1)  $A+B = B+A$
- 2)  $A+(B+C) = (A+B)+C$
- 3)  $A-(B-C) = A-B+C$

EX-2- Find  $A+B$  and  $A-B$  if:-

$$A = \begin{bmatrix} 2 & 1 & 3 \\ 1 & 0 & -2 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & -2 & 2 \\ 2 & 3 & -1 \end{bmatrix}$$

Sol.-

$$A + B = \begin{bmatrix} 2+1 & 1-2 & 3+2 \\ 1+2 & 0+3 & -2-1 \end{bmatrix} = \begin{bmatrix} 3 & -1 & 5 \\ 3 & 3 & -3 \end{bmatrix}$$

$$A - B = \begin{bmatrix} 2-1 & 1-(-2) & 3-2 \\ 1-2 & 0-(+3) & -2-(-1) \end{bmatrix} = \begin{bmatrix} 1 & 3 & 1 \\ -1 & -3 & -1 \end{bmatrix}$$



3. Multiplication by a scalar:- The matrix  $A$  is multiplied by the scalar  $C$  by multiplying each element of  $A$  by  $c$ :-

$$CA = \begin{bmatrix} ca_{11} & ca_{12} & \dots & ca_{1n} \\ ca_{21} & ca_{22} & \dots & ca_{2n} \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ ca_{m1} & ca_{m2} & \dots & ca_{mn} \end{bmatrix}$$

EX-3- Assume  $A = \begin{bmatrix} 3 & 2 & 1 \\ 0 & 5 & -1 \end{bmatrix}$ , find  $3A$ .

Sol.-

$$3A = \begin{bmatrix} 3*3 & 3*2 & 3*1 \\ 3*0 & 3*5 & 3*(-1) \end{bmatrix} = \begin{bmatrix} 9 & 6 & 3 \\ 0 & 15 & -3 \end{bmatrix}$$

4. Matrix multiplication:- For the matrix product  $AB$  to be defined it is necessary that the number of columns of  $A$  be equal to the number of rows of  $B$ . The dimensions of such matrices are said to be conformable. If  $A$  is of dimensions  $m \times p$  and  $B$  is  $p \times n$ , then the  $ij$ th element of the product  $C=AB$  is computed as:-

$$C_{ij} = \sum_{k=1}^p a_{ik} b_{kj}$$

This is the sum of the products of corresponding elements in the  $i$ th row of  $A$  and  $j$ th column of  $B$ . The dimensions of  $AB$  are of course  $m \times n$ .

EX-4- Assume  $A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 1 \end{bmatrix}$  and  $B = \begin{bmatrix} 6 & 5 & 4 \\ -1 & 1 & -1 \\ 0 & 2 & 0 \end{bmatrix}$  find  $AB$ .

Sol.-

$$AB = \begin{bmatrix} 1*6+2(-1)+3*0 & 1*5+2*1+3*2 & 1*4+2(-1)+3*0 \\ -1*6+0(-1)+1*0 & -1*5+0*1+1*2 & -1*4+0(-1)+1*0 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 13 & 2 \\ -6 & -3 & -4 \end{bmatrix}$$



*Properties of multiplication:-*

- a)  $A(B + C) = AB + AC$     *distributive law*
- b)  $A(BC) = (AB)C$     *associative law*
- c)  $AB \neq BA$     *commutative law does not hold*
- d)  $AI = IA = A$

EX-5- Assume  $A = \begin{bmatrix} 1 & 2 \\ 0 & 3 \end{bmatrix}$  and  $B = \begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix}$ , verify that  $AB \neq BA$ .

Sol.-

$$AB = \begin{bmatrix} 1 & 2 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 7 & 1 \\ 6 & 3 \end{bmatrix} \quad \& \quad BA = \begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 2 & 7 \end{bmatrix}$$

*Hence  $AB \neq BA$*

**5. Transpose of matrix:-** Let  $A$  is any  $m \times n$  matrix the transpose of  $A$  is  $n \times m$  matrix  $A'$  formed by interchanging the role of rows and columns.

$$A' = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}' = \begin{bmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & \dots & a_{m2} \\ \dots & \dots & \dots & \dots \\ a_{1n} & a_{2n} & \dots & a_{mn} \end{bmatrix}$$

If a matrix is square and equal to its transpose, it is said to be symmetric, then  $a_{ij} = a_{ji}$  for all pairs of  $i$  and  $j$ .

*Properties of transpose are:-*

- a)  $(A \mp B)' = A' \mp B'$
- b)  $(AB)' = B'A'$
- c)  $(A')' = A$

EX-6- Assume  $A = \begin{bmatrix} 3 & 2 & 5 \\ 2 & -1 & 4 \\ 5 & 4 & 0 \end{bmatrix}$  and  $B = \begin{bmatrix} 4 & -1 & 0 \\ 5 & 4 & 3 \\ 2 & 1 & -1 \end{bmatrix}$ , show that:-

- 1)  $A$  is symmetric matrix
- 2)  $(A + B)' = A' + B'$
- 3)  $(AB)' = B'A'$



Sol.-

$$1) A' = \begin{bmatrix} 3 & 2 & 5 \\ 2 & -1 & 4 \\ 5 & 4 & 0 \end{bmatrix}' = \begin{bmatrix} 3 & 2 & 5 \\ 2 & -1 & 4 \\ 5 & 4 & 0 \end{bmatrix} = A \Rightarrow A \text{ is a symmetric matrix.}$$

$$2) L.H.S. = (A+B)' = \begin{bmatrix} 7 & 1 & 5 \\ 7 & 3 & 7 \\ 7 & 5 & -1 \end{bmatrix}' = \begin{bmatrix} 7 & 7 & 7 \\ 1 & 3 & 5 \\ 5 & 7 & -1 \end{bmatrix}$$

$$R.H.S. = A' + B' = \begin{bmatrix} 3 & 2 & 5 \\ 2 & -1 & 4 \\ 5 & 4 & 0 \end{bmatrix} + \begin{bmatrix} 4 & 5 & 2 \\ -1 & 4 & 1 \\ 0 & 3 & -1 \end{bmatrix} = \begin{bmatrix} 7 & 7 & 7 \\ 1 & 3 & 5 \\ 5 & 7 & -1 \end{bmatrix} = L.H.S.$$

$$\therefore (A+B)' = A' + B'$$

$$3) L.H.S. = (AB)' = \begin{bmatrix} 32 & 10 & 1 \\ 11 & -2 & -7 \\ 40 & 11 & 12 \end{bmatrix}' = \begin{bmatrix} 32 & 11 & 40 \\ 10 & -2 & 11 \\ 1 & -7 & 12 \end{bmatrix}$$

$$R.H.S. = B'A' = \begin{bmatrix} 4 & 5 & 2 \\ -1 & 4 & 1 \\ 0 & 3 & -1 \end{bmatrix} \begin{bmatrix} 3 & 2 & 5 \\ 2 & -1 & 4 \\ 5 & 4 & 0 \end{bmatrix} = \begin{bmatrix} 32 & 11 & 40 \\ 10 & -2 & 11 \\ 1 & -7 & 12 \end{bmatrix} = L.H.S.$$

$$\therefore (AB)' = B'A'$$