

## LEC 8                    Mucogingival surgery

Periodontal treatment involving procedures for correction of defects in morphology, position and/or the amount of soft tissue (gingiva and alveolar mucosa) and underlying bone support at teeth and implants. These procedures are varied from simple gingivectomy or crown lengthening procedures to complex gingival grafting procedures. In patients with bone defects GTR and bone grafting (Guided bone regeneration, GBR) may also be employed to increase the bulk of available alveolar bone.

**Grafting procedures generally aim to cover exposed roots, to increase the bulk of the width of keratinized gingiva and to prevent further gingival recession.**


### **Grafting procedures include**

- Free gingival graft (epithelium + connective tissue) - The pedicle sliding graft



-The sub epithelial connective tissue graft (connective tissue)

-Lateral reposition flap



### **Advantages**

- a. One surgical site
- b. Good vascularity of the pedicle flap.
- c. Ability to cover isolated, denuded roots that have adequate donor tissue laterally.

### **Disadvantages**

- a. Limited by the amount of adjacent keratinized attached gingiva.
- b. Possibility of recession at the donor site.
- c. Dehiscence or fenestration at the donor site.
- d. Limited to one or two teeth with gingival recession.

### **Indications:**

- a. For covering the isolated denuded root.
- b. When there is sufficient width of interdental papilla in the adjacent teeth, and sufficient vestibular depth.

### **Contraindications:**

- a. Presence of deep interproximal pockets.
- b. Excessive root prominence.
- c. Deep or extensive root abrasion or erosion.

## **Frenectomy and frenotomy**

Techniques for the removal of the frenum

A frenum is a fold of mucous membrane, usually with enclosed muscle fibers, that attaches the lips and cheeks to the alveolar mucosa and/or gingiva and underlying periosteum.

**A frenum becomes a problem if the attachment is too close to the marginal gingiva. Tension on the frenum may pull the gingival margin away from the tooth. This condition may be conducive to plaque accumulation and inhibit proper brushing of the teeth with pocket formation. Also may tend to open the sulcus and gingival recession.**

### **Frenectomy or Frenotomy**

The term frenectomy is **complete removal** of the frenum, including its attachment to underlying bone and may be required in the correction of an abnormal diastema between maxillary central incisors. Frenotomy is the incision of the frenum and relocating the frenal attachment.

**Frenal problems occur most often on the facial surface between maxillary and mandibular central incisors and in the canine and premolar areas. They occur less often on the lingual surface of the mandible.**

**The technique for the removal of the frenum accomplished as follows:**

1. After anesthetizing the area, engage the frenum with a hemostat inserted to the depth of the vestibule.

2. Incise along the upper surface of the hemostat, extending beyond the tip.

3. Make a similar incision along the undersurface of the hemostat.

4. Remove the triangular resected portion of the frenum with the hemostat. This exposes the underlying brushlike fibrous attachment to the bone.

5. Make a horizontal incision, separating the fibers, and bluntly dissect to the bone.

6. Undermining the incision to approximate the border of incisions for suturing.

7. Clean the field of operation and pack with gauze sponges until bleeding stops.

8. Cover the area with periodontal pack

9. Remove the pack after 1 week.

**One month is usually required** for the formation of an intact mucosa with the frenum attached in its new position.

#### **Periodontal dressing: are mainly used :**

1. To protect the wound post surgically

2- To obtain and maintain a close adaptation of the mucosal flaps to the underlying bone (especially when a flap has been repositioned apically)

3- For the comfort of the patient

4- Prevent post operative bleeding during the initial phase of healing

5- Prevent the formation of excessive granulation tissue

## **Periodontal dressing should have the following properties:**

- 1-Should be soft but still have enough plasticity and flexibility to facilitate its placement in the operated area and to allow proper adaptation.
- 2-Should harden within a reasonable time
- 3-After setting should be sufficiently rigid to prevent fracture and dislocation.
- 4-Should have a smooth surface after setting to prevent irritation to the cheek and lips
- 5-Should preferably have bactericidal properties to prevent excessive plaque formation
- 6-Not detrimentally interfere with healing

## **Types of dressing**

**1-Zinc-oxide eugenol pack:** eugenol in this type may induce an allergic reaction

**2-Non eugenol pack:** e.g. **Coe pack**; one tube contain zinc oxide and lorothidol (Fungicidal) and the second tube contain non ionizing carboxilic acids and chlorothymol (bacteriostatic agent)

**3-Light cured dressing**

## **Maintenance phase(supportive periodontal therapy SPT)**

Preservation of the periodontal health of the treated patient requires a supportive program that is just as important as the therapy used to treat the periodontal disease. **The maintenance phase of periodontal treatment starts immediately after the completion of phase I therapy.** While the patient is in the maintenance phase, the necessary surgical and restorative procedure are performed. This insures that all areas of the mouth retain the degree of health attained after phase I therapy.

### **The primary goal of maintenance therapy include**

I-Maintenance of oral health

2-Prevention of new infection

3-Prevention of re-infection and disease recurrence

**The time interval between the recall appointments should be based on a periodontal risk assessment**(type and severity of periodontitis, systemic and local risk factors, degree of motivation, compliance, manual dexterity and the patient success to maintain a proper personal oral hygiene standard.

It is important to emphasize that the recall program must be designed to meet the individual need of the patient, some patients should be recall every month while other may have to be checked only once a year.

**Findings from long-term clinical trials have suggested that recall appointments, once every three month is effective in preventing**

**disease recurrence.**

**There are three parts to an SPT appointment:**

- 1.examination
- 2.treatment
- 3.report,clean up and scheduling

**The time required for a recall visit for patients with multiple teeth in both arches is approximately 1 hour**

## **Recurrence of Periodontal Disease**

Occasionally, lesions may recur, which is often due to **inadequate** plaque/biofilm control on the part of the patient or **failure** to comply with recommended **SPT** schedules. **It should be understood, however, that it is the responsibility of the dentist to educate and motivate patients to improve their oral hygiene techniques.** Surgery **should not** be undertaken unless the patient participates in disease prevention and demonstrates proficiency in plaque/biofilm control.

**Other causes for recurrence include the following:**

- 1.Inadequate or insufficient treatment that has failed to remove all of the potential factors favoring biofilm accumulation. **Incomplete calculus removal** in areas of difficult access is a **common source** of problems.
2. Inadequate restorations placed after the periodontal treatment was completed.
3. Failure of the patient to return for periodic maintenance care . This may be a result of the patient's conscious or unconscious decision not

to continue treatment or the failure of the dentist and staff to emphasize the need for periodic supportive therapy.


4. Presence of some systemic diseases that may affect host resistance to previously acceptable levels of biofilm.

**A failing case can be recognized by the following:**


1. Recurring inflammation revealed by gingival changes and bleeding of the sulcus on probing.
2. Increasing depth of sulci, leading to the recurrence of pocket formation.
3. Gradual increases in bone loss, as determined by radiographs.
4. Gradual increases in tooth mobility, as ascertained by clinical examination.

The decision to retreat a periodontal patient should not be made at the preventive maintenance appointment but should be postponed for **1 to 2 weeks**. Often, the mouth appears improved at that time because of the resolution of edema and the resulting improved tone of the gingiva.

## FRENECTOMY



## BON GRAFT



## Periodontal dressing