


## Cell Wall Inhibitors

The bacterial cell wall is a rigid outer layer that completely surrounds the cytoplasmic membrane (the mammalian cells do not possess cell wall). Cell wall inhibitors selectively interfere with synthesis of the bacterial cell wall. The cell wall is composed of a polymer called peptidoglycan that consists of glycan units joined to each other by peptide cross-links. Inhibitors of cell wall synthesis require actively proliferating microorganisms.



### I. Penicillins:

Penicillins act by inhibiting the enzymes involved in the cross linking of the peptidoglycan layer of the cell wall, which is weakened, and this leads to osmotic rupture. Penicillins are thus bactericidal and are ineffective against resting organisms which are not making new cell wall. (Bacteriostatics should not be taken with penicillins?)

#### • Natural penicillins:

##### Penicillin G (benzylpenicillin):

- ✓ It is destroyed by gastric acid (acid labile); hence penicillin G is usually given by IV route. It can also be administered by IM route but is painful.
- ✓ Penicillin G is widely distributed in body tissues, but poorly crosses the BBB; although during meningitis, adequate amount reaches the CSF. Penicillin G is rapidly excreted in urine mainly by active tubular secretion.
- ✓ Since renal function is not completely developed in infants and neonates, hence excretion of penicillins is slow.
- ✓ The action of penicillins can be augmented and prolonged by giving probenecid simultaneously
- ✓ The duration of action of penicillin G is increased by combining it with poorly water-soluble compounds, such as procaine (procaine penicillin G) or benzathine (benzathine penicillin G) to yield aqueous suspensions. They are called depot penicillins.

- ✓ Penicillin remains the drug of choice for the treatment of gas gangrene (Clostridium perfringens) and syphilis (Treponema pallidum).

### Penicillin V:

- ✓ Only available in oral formulation,
- ✓ It has a spectrum similar to that of penicillin G, but it is not used for treatment of severe infections because of its limited oral absorption.
- ✓ Penicillin V is more acid stable than is penicillin G and is the oral agent employed in the treatment of less severe infections.

- **Semisynthetic penicillins (aminopenicillins) or (extended spectrum penicillins)**

They have broad antibacterial spectrum and are effective against both gram positive and gram negative organisms. They are hydrolysed by penicillinase.

Ampicillin and amoxicillin These extended-spectrum agents are also widely used in the treatment of respiratory infections, and amoxicillin is employed prophylactically by dentists in high-risk patients for the prevention of bacterial endocarditis.

These drugs are coformulated with  $\beta$ -lactamase inhibitors, such as clavulanic acid or sulbactam, to combat infections caused by  $\beta$ -lactamase-producing organisms.

- **Penicillinase-resistant penicillins (Antistaphylococcal penicillins):**

- ✓ These penicillins are useful for the treatment of infections involving penicillinase-producing bacteria, such as Staphylococcus aureus or S. epidermidis.
- ✓ Methicillin, is no longer used clinically because of its toxicity (interstitial nephritis).
- ✓ The penicillinase-resistant penicillins have minimal to no activity against gram-negative infections.

- **Antipseudomonal penicillin:**

- ✓ Piperacillin is also referred to as an antipseudomonal penicillin because of its activity against Pseudomonas aeruginosa.
- ✓ Formulation of piperacillin with tazobactam (available only as parenteral preparation) extends the antimicrobial spectrum to include penicillinase-producing organisms.

### #Adverse reactions to penicillins:

1. Hypersensitivity reactions range from rashes to angioedema (marked swelling of the lips, tongue, and periorbital area) and anaphylaxis. Cross-allergic reactions occur among the  $\beta$ lactam antibiotics. To determine whether treatment with a  $\beta$ -lactam is safe when an allergy is noted, patient history regarding severity of previous reaction is essential.



2. Diarrhea is a common problem that is caused by a disruption of the normal balance of intestinal microorganisms. It occurs to a greater extent with those agents that are incompletely absorbed and have an extended antibacterial spectrum. Pseudomembranous colitis from Clostridium difficile and other organisms may occur with penicillin use.

3. Nephritis: Penicillins, particularly methicillin, have the potential to cause acute interstitial nephritis. [Note: Methicillin is therefore no longer used clinically].

4. Neurotoxicity: The penicillins are irritating to neuronal tissue, and they can provoke seizures if injected intrathecally or if very high blood levels are reached.

5. Hematologic toxicities: Decreased coagulation may be observed with high doses of piperacillin and nafcillin (and, to some extent, with penicillin G). Cytopenias have been associated with therapy of greater than 2 weeks, and therefore, blood counts should be monitored weekly for such patients.

## II. Cephalosporins:

The cephalosporins are  $\beta$ -lactam antibiotics closely related both structurally and functionally to penicillins. They tend to be more resistant than the penicillins to certain  $\beta$ -lactamases.

Cephalosporins have been classified as first, second, third, fourth, and advanced generation, based largely on their bacterial susceptibility patterns and resistance to  $\beta$ -lactamases.

1. First generation: cefazolin and oral cephalexin act as penicillin G substitutes. They are resistant to the staphylococcal Penicillinase. They are the most active cephalosporins for Gram-positive bacterial infections

2. Second generation: cefroxitin and oral cefuroxime display greater activity against gram-negative organisms.

An important use of first- and second-generation cephalosporins is prophylaxis during surgery if an infection is likely to occur.

3. Third generation: These cephalosporins have assumed an important role in the treatment of infectious diseases. Ceftriaxone and cefotaxime have become agents of choice in the treatment of meningitis. Cefixime (Suprax) is an oral preparation.


4. Fourth generation: Cefepime must be administered parenterally. Cefepime has a wide antibacterial spectrum.

5. Advanced generation: Ceftaroline is a broad-spectrum, advanced-generation cephalosporin. It is the only  $\beta$ -lactam with activity against MRSA, and it is indicated for the treatment of complicated skin and skin structure infections and community-acquired pneumonia.

The twice-daily dosing regimen also limits use outside of an institutional setting.

1 <sup>st</sup> GENERATION	2 <sup>nd</sup> GENERATION	3 <sup>rd</sup> GENERATION	4 <sup>th</sup> GENERATION	5 <sup>th</sup> GENERATION
<ul style="list-style-type: none"> <li>• CEFAZOLIN</li> <li>• CEFALOTHIN</li> <li>• CEFALORIDINE</li> <li>• CEFADROXIL</li> <li>• CEPHALEXIN</li> <li>• CEPHRADINE</li> </ul>	<ul style="list-style-type: none"> <li>• CEFOXITIN</li> <li>• CEFPROZIL</li> <li>• CEFOTITAN</li> <li>• CEFMETAZOLE</li> <li>• CEFACLOR</li> <li>• CEFUROXIME</li> </ul>	<ul style="list-style-type: none"> <li>• CEFEXIME</li> <li>• CEFTAZIDIME</li> <li>• CEFOTAXIME</li> <li>• CEFTIZOXIME</li> <li>• CEFPODOXIME</li> <li>• CEFTRIAZONE</li> <li>• CEFOPERAZONE</li> <li>• CEFTIBUTEN</li> <li>• CEFDINIR</li> <li>• MOXALACTAM</li> </ul>	<ul style="list-style-type: none"> <li>• CEFEPIME</li> <li>• CEFPIROME</li> </ul>	<ul style="list-style-type: none"> <li>• CEFTOBIPROLE</li> <li>• CEFTAROLINE</li> </ul>

- Drugs that have 'FA' or 'PHA/PHRA' in their name are 1<sup>st</sup> generation except Cefaclor which is 2<sup>nd</sup> generation drug
- Drugs that end with 'IME' or 'ONE' or 'TEN' are 3<sup>rd</sup> generation except Cefuroxime which is 2<sup>nd</sup> generation drug
- Drugs that have 'PI' in their name are 4<sup>th</sup> generation
- Drugs that have 'ROL' in their name are 5<sup>th</sup> generation



### #Adverse effects:

Like the penicillins, the cephalosporins are generally well tolerated. However, allergic reactions are a concern.

Patients who have had an anaphylactic response, Stevens-Johnson syndrome, or toxic epidermal necrolysis to penicillins should not receive cephalosporins. Cephalosporins should be avoided or used with caution in individuals with penicillin allergy.

The highest rate of allergic cross-sensitivity is between penicillin and first-generation cephalosporins.

### III. Other $\beta$ -Lactam Antibiotics:

- **Carbapenems:** are synthetic  $\beta$ -lactam antibiotics that inhibit cell wall transpeptidation. This results in bactericidal activity against most bacteria. *Imipenem*, *meropenem*, *doripenem*, and *ertapenem* are drugs in this group. Imipenem undergoes cleavage by a dehydropeptidase found in the brush border of the proximal renal tubule. Compounding imipenem with *cilastatin* protects the parent drug from renal dehydropeptidase and, thus, prolongs its activity in the body. The other carbapenems do not require coadministration of cilastatin.



- **Monobactams:** *Aztreonam*

1. It decreases cell wall formation and thus is bactericidal.
2. Only aerobic Gram-negative bacteria, especially *Pseudomonas*, are affected. There is no activity against Gram-positive bacteria or anaerobes.
3. It is resistant to most  $\beta$ -lactamases.
4. Kinetics are similar to the penicillins, although it must be administered parenterally.
5. There is no cross-allergenicity with penicillins.

### IV. $\beta$ -Lactamase Inhibitors:

$\beta$ -Lactamase inhibitors, such as clavulanic acid, sulbactam, and tazobactam, contain a  $\beta$ -lactam ring but, by themselves, do not have significant antibacterial activity or cause any significant adverse effects. They function by inactivating  $\beta$ -lactamases, thereby protecting the antibiotics that are normally substrates for these enzymes. The  $\beta$ -lactamase inhibitors are, therefore, formulated in combination with  $\beta$ -lactamase-sensitive antibiotics, such as amoxicillin, ampicillin, and piperacillin.



## V. Vancomycin:

Following cell entry, it binds to peptidoglycan precursors, disrupting polymerization and cross-linking required for maintenance of cell wall integrity. This interaction results in bactericidal activity.

Due to an increase in MRSA, vancomycin is commonly used in patients with skin and soft tissue infections, infective endocarditis, and nosocomial pneumonia (hospital acquired pneumonia).


Common adverse events include nephrotoxicity, infusion-related reactions (red man syndrome and phlebitis), and ototoxicity.

Vancomycin has poor absorption after oral administration, so use of the oral formulation is limited to the management of *Clostridium difficile* infection in the colon.



## VI. Daptomycin:

Daptomycin is a bactericidal antibiotic. It is indicated for the treatment of complicated skin and skin structure infections and bacteremia caused by *S. aureus*, including those with right-sided infective endocarditis. Efficacy of treatment with daptomycin in left-sided endocarditis has not been demonstrated. Additionally, daptomycin is inactivated by pulmonary surfactants; thus, it should never be used in the treatment of pneumonia. Daptomycin is dosed IV once daily.



## VII. Fosfomycin

Fosfomycin is a bactericidal synthetic derivative of phosphonic acid. It blocks cell wall synthesis. It is indicated for urinary tract infections caused by *E. coli* or *E. faecalis* and is considered first-line therapy for acute cystitis. Due to its unique structure and mechanism of action, cross-resistance with other antimicrobial agents is unlikely. Fosfomycin is rapidly absorbed after oral administration and distributes well to the kidneys, bladder, and prostate. The drug is excreted in its active form in the urine and maintains high concentrations over several days, allowing for a one-time dose.

The most commonly reported adverse effects include diarrhea, vaginitis, nausea, and headache.



## VIII. Polymyxins

The polymyxins bind to phospholipids on the bacterial cell membrane of gram-negative bacteria. They have a detergent-like effect that disrupts cell membrane integrity, leading to leakage of cellular components and cell death.

Only two forms of polymyxin are in clinical use today, polymyxin B and colistin (polymyxin E).

The use of these drugs has been limited due to the increased risk of nephrotoxicity and neurotoxicity when used systemically. However, with increasing gram-negative resistance, they are now commonly used as salvage therapy for patients with multidrug-resistant infections.

